

2-D SKEW CONSTACYCLIC CODES OVER $R[x, y; \rho, \theta]$

H. MOSTAFANASAB

ABSTRACT. For a finite field \mathbb{F}_q , the bivariate skew polynomial ring $\mathbb{F}_q[x, y; \rho, \theta]$ has been used to study codes [25]. In this paper, we give some characterizations of the ring $R[x, y; \rho, \theta]$, where R is a commutative ring. We investigate 2-D skew (λ_1, λ_2) -constacyclic codes in the ring $R[x, y; \rho, \theta]/\langle x^l - \lambda_1, y^s - \lambda_2 \rangle_l$. Also, the dual of 2-D skew (λ_1, λ_2) -constacyclic codes is investigated.

1. INTRODUCTION

Cyclic codes are amongst the most studied algebraic codes. Their structure is well known over finite fields [17]. Recently codes over rings have generated a lot of interest after a breakthrough paper by Hammons et al. [12] showed that some well known binary non-linear codes are actually images of some linear codes over \mathbb{Z}_4 under the Gray map. Constacyclic codes over finite fields form a remarkable class of linear codes, as they include the important family of cyclic codes. Constacyclic codes also have practical applications as they can be efficiently encoded using simple shift registers. They have rich algebraic structures for efficient error detection and correction, which explains their preferred role in engineering. In general, due to their rich algebraic structure, constacyclic codes have been studied over various finite chain rings (see [1, 2, 7, 8, 9, 10, 18, 24]). Two-dimensional (2-D) cyclic codes are generalizations of usual cyclic codes which were introduced by Ikai et al. [13] and Imai [14]. Güneri and Özbudak studied the relations

MSC(2010): Primary: 00A69; Secondary: 12E20; 94B05

Keywords: cyclic codes, skew polynomial rings, 2-D skew constacyclic codes.

Received: 26 May 2016, Accepted: 19 December 2016.

between quasi-cyclic codes and 2-D cyclic codes [11]. Decoding problem for 2-D cyclic codes was studied by some authors [20, 21, 22, 23]. Polynomial rings and their ideals are essential to the construction and understanding of cyclic codes. For the first time in [3] non-commutative skew polynomial rings have been used to construct (a generalization of) cyclic codes. Skew-cyclic codes were introduced by Boucher et al. [5, 6]. They considered the skew-cyclic codes as ideals or submodules over skew polynomial rings and studied dual skew-cyclic codes. Skew constacyclic codes have been investigated by Boucher et al. in [4] and Jitman et al. in [15]. Xiuli and Hongyan [25] generalized the 2-D cyclic codes to 2-D skew-cyclic codes. They studied the structures and properties of 2-D skew-cyclic codes. Also, they built relationships between 2-D skew-cyclic codes and other known codes.

Throughout this paper, let R be a commutative ring. For two given automorphisms ρ and θ of R , we consider the set of formal bivariate polynomials

$$R[x, y; \rho, \theta] = \left\{ \sum_{j=0}^t \sum_{i=0}^k a_{i,j} x^i y^j \mid a_{i,j} \in R \text{ and } k, t \in \mathbb{N}_0 \right\}$$

which forms a ring under the usual addition of polynomials and where the multiplication is defined using the rule

$$ax^i y^j \star bx^r y^s = a\rho^i \theta^j(b) x^{i+r} y^{j+s},$$

and extended to all elements of $R[x, y; \rho, \theta]$ by associativity and distributivity. The ring $R[x, y; \rho, \theta]$ is called a *bivariate skew polynomial ring over R* and an element in $R[x, y; \rho, \theta]$ is called a *bivariate skew polynomial*. It is easy to see that $R[x, y; \rho, \theta]$ is a non-commutative ring unless ρ and θ are identity automorphisms on R . For a bivariate skew polynomial $f(x, y)$ in $R[x, y; \rho, \theta]$, let $\langle f(x, y) \rangle_l$ denote the left ideal of $R[x, y; \rho, \theta]$ generated by $f(x, y)$. Note that $\langle f(x, y) \rangle_l$ does not need to be two-sided.

In section 2, we give some characterizations of the ring $R[x, y; \rho, \theta]$. In section 3, we introduce and investigate 2-D skew (λ_1, λ_2) -constacyclic codes in $R[x, y; \rho, \theta]/\langle x^l - \lambda_1, y^s - \lambda_2 \rangle_l$. Also, the dual of 2-D skew (λ_1, λ_2) -constacyclic codes is investigated.

Let $\mathbb{N}_0^2 = \mathbb{N}_0 \times \mathbb{N}_0$. Then \mathbb{N}_0^2 is a partial ordered set with $(i, j) \geq (k, l)$ if and only if $i \geq k$ and $j \geq l$. Moreover, \mathbb{N}_0^2 can be also totally ordered by a kind of lexicographic order “ \Rightarrow ”, where $(i, j) \Rightarrow (k, l)$ if and only if $j > l$ or both $j = l$ and $i \geq k$. Otherwise, $(i, j) \rightarrow (k, l)$ means $j > l$ or both $j = l$ and $i > k$. Notice that $(i, j) \geq (k, l)$ implies that $(i, j) \Rightarrow (k, l)$, but the converse does not necessarily hold. A nonzero

bivariate polynomial $f(x, y) \in R[x, y; \rho, \theta]$ is said to have quasi-degree $\deg(f(x, y)) = (k, t)$ if $f(x, y)$ has a nonzero term $a_{k,t}x^k y^t$ but does not have any nonzero term $a_{i,j}x^i y^j$ such that $(i, j) \rightarrow (k, t)$ holds. In this case, $a_{k,t}$ is called *the leading coefficient of $f(x, y)$* . A bivariate skew polynomial is called *monic* provided its leading coefficient is 1. Let \mathbb{F}_q be the Galois field with q elements. For any polynomials $f(x, y)$ and $g(x, y)$ in $\mathbb{F}_q[x, y; \rho, \theta]$ we have

$$\deg(f(x, y) \star g(x, y)) = \deg(f(x, y)) + \deg(g(x, y)).$$

It is straightforward to see that $\mathbb{F}_q[x, y; \rho, \theta]$ has no nonzero zero-divisors.

2. BASIC PROPERTIES OF $R[x, y; \rho, \theta]$

In this paper we denote by $R^{\rho, \theta}$ (resp. $R^{\rho\theta}$) the subring of R that is fixed by ρ, θ (resp. $\rho\theta$).

Let R be a commutative ring and $f(x) \in R[x]$. In [16], McCoy observed that if $0 \neq g(x) \in R[x]$ be such that $f(x)g(x) = 0$, then there exists a nonzero element r of R such that $f(x)r = 0$. Now, we state the bivariate skew version of the McCoy condition.

Theorem 2.1. *Let $f(x, y) \in R^{\rho, \theta}[x, y; \rho, \theta]$. If $f(x, y) \star g(x, y) = 0$ for some $0 \neq g(x, y) \in R[x, y; \rho, \theta]$, then there exists $0 \neq r \in R$ such that $f(x, y) \star r = 0$.*

Proof. Suppose that $f(x, y) = \sum_{(i,j) \in \Lambda} f_{i,j}x^i y^j$. We can assume that

$g(x, y)$ is of the minimal degree $\deg(g(x, y)) = (u, v) (\rightarrow (0, 0))$ (with respect to “ \Rightarrow ”), and $g(x, y)$ has the leading coefficient $g_{u,v}$. By the minimality of $\deg(g(x, y))$ we have $f(x, y) \star g_{u,v} \neq 0$. If $f_{i,j}g_{u,v} = 0$ for every $(i, j) \in \Lambda$, then $f_{i,j}g_{u,v} = 0$ for every $(i, j) \in \Lambda$. Now, since $f(x, y) \in R^{\rho, \theta}[x, y; \rho, \theta]$, then $f(x, y) \star g_{u,v} = 0$ which is a contradiction. Therefore $f_{i,j}g_{u,v} \neq 0$ for some $(i, j) \in \Lambda$ and so assume that (k, t) is the largest pair with this property (with respect to

“ \Rightarrow ”). Hence $0 = f(x, y) \star g(x, y) = \left(\sum_{j=0}^t \sum_{i=0}^k f_{i,j}x^i y^j \right) \star g(x, y)$ which

implies that $\rho^i \theta^j (f_{k,t} g_{u,v}) = f_{k,t} \rho^i \theta^j (g_{u,v}) = 0$. Then $f_{k,t} g_{u,v} = 0$. Thus $\deg(g(x, y)) \rightarrow \deg(f_{k,t} g(x, y))$. On the other hand $f(x, y) \star (f_{k,t} g(x, y)) = f(x, y) \star (g(x, y) \star f_{k,t}) = 0$ which contradicts the minimality of $\deg(g(x, y))$. \square

The center of a ring S , denoted by $Z(S)$, is the subset of S consisting of all those elements in S that commute with every element in S .

Proposition 2.2. *Let $g(x, y) \in Z(R[x, y; \rho, \theta])$. Then $\langle x^k y^t \star g(x, y) \rangle_l$ is a two-sided ideal in $R[x, y; \rho, \theta]$ for every $k, t \in \mathbb{N}_0$.*

Proof. Let $g(x, y) \in Z(R[x, y; \rho, \theta])$. Assume that $f(x, y) = \sum_{j=0}^n \sum_{i=0}^m a_{i,j} x^i y^j \in R[x, y; \rho, \theta]$. Then

$$\begin{aligned}
(x^k y^t \star g(x, y)) \star f(x, y) &= (x^k y^t \star g(x, y)) \star \sum_{j=0}^n \sum_{i=0}^m a_{i,j} x^i y^j \\
&= \left(x^k y^t \star \left(\sum_{j=0}^n \sum_{i=0}^m a_{i,j} x^i y^j \right) \right) \star g(x, y) \\
&\quad (\text{since } g(x, y) \text{ is central}) \\
&= \left(\sum_{j=0}^n \sum_{i=0}^m \rho^k \theta^t (a_{i,j}) x^{k+i} y^{t+j} \right) \star g(x, y) \\
&= \left(\left(\sum_{j=0}^n \sum_{i=0}^m \rho^k \theta^t (a_{i,j}) x^i y^j \right) \star x^k y^t \right) \star g(x, y) \\
&= \left(\sum_{j=0}^n \sum_{i=0}^m \rho^k \theta^t (a_{i,j}) x^i y^j \right) \star (x^k y^t \star g(x, y)),
\end{aligned}$$

which belongs to $\langle x^k y^t g(x, y) \rangle_l$. Hence, the result follows. \square

Proposition 2.3. *Let $|\langle \rho \rangle| \mid l$ and $|\langle \theta \rangle| \mid s$. Then $R^{\rho, \theta}[x^l, y^s; \rho, \theta] \subseteq Z(R[x, y; \rho, \theta])$.*

Proof. Let $a \in R^{\rho, \theta}$, $b \in R$ and $i, j, k, t \in \mathbb{N}_0$. Therefore

$$\begin{aligned}
(ax^{li} y^{sj}) \star (bx^k y^t) &= a \rho^{li} \theta^{sj} (b) x^{li+k} y^{sj+t} \\
&= b a x^{k+li} y^{t+sj} \\
&= (bx^k y^t) \star (\theta^{-t} \rho^{-k} (a) x^{li} y^{sj}) \\
&= (bx^k y^t) \star (ax^{li} y^{sj}) \quad (\text{since } a \in R^{\rho^{-1}, \theta^{-1}}).
\end{aligned}$$

Consequently $R^{\rho, \theta}[x^l, y^s; \rho, \theta] \subseteq Z(R[x, y; \rho, \theta])$. \square

Proposition 2.4. *Let λ be a unit in R . The following conditions hold:*

- (1) *If $\langle x^l - \lambda \rangle_l$ is a two-sided ideal of $R[x, y; \rho, \theta]$, then $|\langle \rho \rangle| \mid l$ and $\rho(\lambda) = \lambda$.*
- (2) *If $\langle y^s - \lambda \rangle_l$ is a two-sided ideal of $R[x, y; \rho, \theta]$, then $|\langle \theta \rangle| \mid s$ and $\theta(\lambda) = \lambda$.*

Proof. The proof is similar to that of [15, Proposition 2.2]. \square

Proposition 2.5. *Let λ_1, λ_2 be units in R . The following conditions hold:*

- (1) *If $|\langle \theta \rangle| \mid s$ and $\langle x^l - \lambda_1, y^s - \lambda_2 \rangle_l$ is a two-sided ideal of $R[x, y; \rho, \theta]$, then $|\langle \rho \rangle| \mid l$ and $\rho(\lambda_1) = \lambda_1$.*
- (2) *If $|\langle \rho \rangle| \mid l$ and $\langle x^l - \lambda_1, y^s - \lambda_2 \rangle_l$ is a two-sided ideal of $R[x, y; \rho, \theta]$, then $|\langle \theta \rangle| \mid s$ and $\theta(\lambda_2) = \lambda_2$.*

Proof. Suppose that $|\langle \theta \rangle| \mid s$ and $\langle x^l - \lambda_1, y^s - \lambda_2 \rangle_l$ is a two-sided ideal. Let $r \in R$. Then $rx^l - r\lambda_1 = r(x^l - \lambda_1) = (x^l - \lambda_1)r'$ for some $r' \in R$. The remainder is similar to the proof of [15, Proposition 2.2]. \square

Theorem 2.6. *Let $\lambda \in R$. The following conditions hold:*

- (1) *If $|\langle \rho \rangle| \mid l$ and $\lambda \in R^{\rho, \theta}$, then $x^l - \lambda$ is central in $R[x, y; \rho, \theta]$.*
- (2) *If $|\langle \theta \rangle| \mid s$ and $\lambda \in R^{\rho, \theta}$, then $y^s - \lambda$ is central in $R[x, y; \rho, \theta]$.*
- (3) *If $|\langle \rho \rangle| \mid l$, $|\langle \theta \rangle| \mid s$ and $\lambda_1, \lambda_2 \in R^{\rho, \theta}$, then $\langle x^l - \lambda_1, y^s - \lambda_2 \rangle_l$ is a two-sided ideal of $R[x, y; \rho, \theta]$.*

Proof. (1) Suppose that $|\langle \rho \rangle| \mid l$, $\rho(\lambda) = \lambda$ and $\theta(\lambda) = \lambda$. Let $a \in R$ and $k, t \in \mathbb{N}_0$. Thus

$$\begin{aligned}
(ax^k y^t) \star (x^l - \lambda) &= ax^{k+l} y^t - ax^k y^t \lambda \\
&= ax^{l+k} y^t - \rho^k \theta^t(\lambda) ax^k y^t \\
&= (x^l \star \rho^{-l}(a) x^k y^t) - \lambda ax^k y^t \\
&= (x^l \star ax^k y^t) - \lambda ax^k y^t \\
&= (x^l - \lambda) \star (ax^k y^t).
\end{aligned}$$

Consequently $x^l - \lambda \in Z(R[x, y; \rho, \theta])$.

(2) The proof is similar to that of (1).

(3) Is a direct consequence of parts (1) and (2). \square

Corollary 2.7. *Let R be a ring. The following conditions hold:*

- (1) *Let $\theta(\lambda) = \lambda$. Then $|\langle \rho \rangle| \mid l$ and $\rho(\lambda) = \lambda$ if and only if $x^l - \lambda$ is central in $R[x, y; \rho, \theta]$.*
- (2) *Let $\rho(\lambda) = \lambda$. Then $|\langle \theta \rangle| \mid s$ and $\theta(\lambda) = \lambda$ if and only if $y^s - \lambda$ is central in $R[x, y; \rho, \theta]$.*

Proof. By Proposition 2.4 and Theorem 2.6. \square

Proposition 2.8. *Let $f \star g$ be a monic central bivariat skew polynomial in $R[x, y; \rho, \theta]$ for some $f, g \in R[x, y; \rho, \theta]$. Then $f \star g = g \star f$.*

Proof. Since $f \star g$ is central, then we have $(g \star f) \star g = g \star (f \star g) = (f \star g) \star g$. Therefore $(g \star f - f \star g) \star g = 0$. On the other hand $f \star g$ is monic, so the leading coefficient of g is unit. Hence g is not a zero-divisor. Consequently $f \star g = g \star f$ in $R[x, y; \rho, \theta]$. \square

Corollary 2.9. *Let $|\langle \rho \rangle| \mid l$, $|\langle \theta \rangle| \mid s$ and $\lambda_1, \lambda_2 \in R^{\rho, \theta}$. If $(x^l - \lambda_1) \star (y^s - \lambda_2) = f \star g$ for some $f, g \in R[x, y; \rho, \theta]$, then $f \star g = g \star f$.*

Proof. By Theorem 2.6 and Proposition 2.8. □

3. 2-D SKEW (λ_1, λ_2) -CONSTACYCLIC CODES

Definition 3.1. Let \mathcal{C} be a linear code over R of length ls whose codewords are viewed as $l \times s$ arrays, i.e., $c \in \mathcal{C}$ is written as

$$c = \begin{pmatrix} c_{0,0} & c_{0,1} & \cdots & c_{0,s-1} \\ c_{1,0} & c_{1,1} & \cdots & c_{1,s-1} \\ \vdots & \vdots & \ddots & \vdots \\ c_{l-1,0} & c_{l-1,1} & \cdots & c_{l-1,s-1} \end{pmatrix}.$$

We say that \mathcal{C} is a *column skew λ_1 -constacyclic code* if for every $l \times s$ array $c = (c_{i,j}) \in \mathcal{C}$ we have that

$$\begin{pmatrix} \lambda_1 \rho(c_{l-1,0}) & \lambda_1 \rho(c_{l-1,1}) & \cdots & \lambda_1 \rho(c_{l-1,s-1}) \\ \rho(c_{0,0}) & \rho(c_{0,1}) & \cdots & \rho(c_{0,s-1}) \\ \vdots & \vdots & \ddots & \vdots \\ \rho(c_{l-2,0}) & \rho(c_{l-2,1}) & \cdots & \rho(c_{l-2,s-1}) \end{pmatrix} \in \mathcal{C}.$$

Also, we say that \mathcal{C} is a *row skew λ_2 -constacyclic code* if for every $l \times s$ array $c = (c_{i,j}) \in \mathcal{C}$ we have that

$$\begin{pmatrix} \lambda_2 \theta(c_{0,s-1}) & \theta(c_{0,0}) & \cdots & \theta(c_{0,s-2}) \\ \lambda_2 \theta(c_{1,s-1}) & \theta(c_{1,0}) & \cdots & \theta(c_{1,s-2}) \\ \vdots & \vdots & \ddots & \vdots \\ \lambda_2 \theta(c_{l-1,s-1}) & \theta(c_{l-1,0}) & \cdots & \theta(c_{l-1,s-2}) \end{pmatrix} \in \mathcal{C}.$$

If \mathcal{C} is both column skew λ_1 -constacyclic and row skew λ_2 -constacyclic, then we call \mathcal{C} a *2-D skew (λ_1, λ_2) -constacyclic code*.

Define $R^\circ := R[x, y; \rho, \theta] / \langle x^l - \lambda_1, y^s - \lambda_2 \rangle_l$. Consider the R -module isomorphism $R^{l \times s} \rightarrow R^\circ$ defined by $(a_{i,j}) \mapsto \sum_{j=0}^{s-1} \sum_{i=0}^{l-1} a_{i,j} x^i y^j$ where $R^{l \times s}$ denotes the set of all $l \times s$ arrays. Then a codeword $c \in \mathcal{C}$ can be denoted by bivariate skew polynomial $c(x, y)$ under above isomorphism.

Theorem 3.2. *A code \mathcal{C} in R° is a 2-D skew (λ_1, λ_2) -constacyclic code if and only if \mathcal{C} is a left $R[x, y; \rho, \theta]$ -submodule of the left $R[x, y; \rho, \theta]$ -module R° .*

Proof. Similar to the proof of [25, Theorem 3.2]. □

As a direct consequence of Theorem 2.6(3) and Theorem 3.2 we have the next result.

Corollary 3.3. *Let $|\langle \rho \rangle| \mid l$, $|\langle \theta \rangle| \mid s$ and $\lambda_1, \lambda_2 \in R^{\rho, \theta}$. A code \mathcal{C} in R° is a 2-D skew (λ_1, λ_2) -constacyclic code if and only if \mathcal{C} is a left ideal of R° .*

Theorem 3.4. *Let $f_1(x, y), f_2(x, y) \in R[x, y; \rho, \theta]$ be two nonzero bivariate polynomials where $f_2(x, y)$ is monic. Provided that $\deg(f_1(x, y)) \geq \deg(f_2(x, y))$, there exists a pair of polynomials $h(x, y) (\neq 0), g(x, y) \in R[x, y; \rho, \theta]$ which satisfy $f_1(x, y) = h(x, y) \star f_2(x, y) + g(x, y)$ such that $g(x, y) = 0$ or $\deg(f_2(x, y)) \not\leq \deg(g(x, y))$ ($\leftarrow \deg(f_1(x, y))$).*

Proof. The proof is similar to that of [25, Theorem 2.1]. \square

Lemma 3.5. *Let \mathcal{C} be a 2-D skew (λ_1, λ_2) -constacyclic code in R° and $g(x, y)$ be a monic polynomial in $R[x, y; \rho, \theta]$. Then $g(x, y)$ is of the minimum degree (with respect to “ \leq ”) in \mathcal{C} if and only if $\mathcal{C} = \langle g(x, y) \rangle_l$.*

Proof. The “if” part is obvious. Let $g(x, y)$ be a monic polynomial of the minimum degree (with respect to “ \leq ”) in \mathcal{C} . If $c(x, y) \in \mathcal{C}$, then by the Division Algorithm in $R[x, y; \rho, \theta]$, there exists a pair of polynomials $h(x, y) (\neq 0), r(x, y) \in R[x, y; \rho, \theta]$ which satisfy $c(x, y) = h(x, y) \star g(x, y) + r(x, y)$, where either $r(x, y) = 0$ or $\deg(g(x, y)) \not\leq \deg(r(x, y))$ ($\leftarrow \deg(c(x, y))$). As \mathcal{C} is an $R[x, y; \rho, \theta]$ -module, $r(x, y) \in \mathcal{C}$ and the minimality of the degree of $g(x, y)$ implies $r(x, y) = 0$. \square

Theorem 3.6. *Let \mathcal{C} be a 2-D skew (λ_1, λ_2) -constacyclic code in \mathbb{F}_q° and $g(x, y)$ be a monic generator polynomial of \mathcal{C} in $\mathbb{F}_q[x, y; \rho, \theta]$. Then the following conditions hold:*

- (1) $g(x, y)$ divides $(x^l - \lambda_1) \star (y^s - \lambda_2)$.
- (2) Suppose that $\deg(g(x, y)) = (l - k, s - t)$. Then

$$\mathbb{B} = \left\{ \begin{array}{cccc} g(x, y), & y \star g(x, y), & \dots & , y^{t-1} \star g(x, y) \\ x \star g(x, y), & xy \star g(x, y), & \dots & , xy^{t-1} \star g(x, y) \\ \vdots & \vdots & \ddots & \vdots \\ x^{k-1} \star g(x, y), & x^{k-1}y \star g(x, y), & \dots & , x^{k-1}y^{t-1} \star g(x, y) \end{array} \right\}$$

is a basis for \mathcal{C} over \mathbb{F}_q and so $|\mathcal{C}| = q^{kt}$.

Proof. (1) Assume that $g(x, y)$ does not divide $(x^l - \lambda_1) \star (y^s - \lambda_2)$. By the Division Algorithm, there exist $f(x, y) (\neq 0), r(x, y) \in \mathbb{F}_q[x, y; \rho, \theta]$ which satisfy

$$(x^l - \lambda_1) \star (y^s - \lambda_2) = f(x, y) \star g(x, y) + r(x, y)$$

such that either $r(x, y) = 0$ or $\deg(g(x, y)) \not\leq \deg(r(x, y))$. If $\deg(g(x, y)) \not\leq \deg(r(x, y))$, then

$$\begin{aligned} r(x, y) &= (x^l - \lambda_1) \star (y^s - \lambda_2) - f(x, y) \star g(x, y) \\ &\equiv -f(x, y) \star g(x, y) \pmod{\langle x^l - \lambda_1, y^s - \lambda_2 \rangle_l} \end{aligned}$$

is also in \mathcal{C} which contradicts the minimality of $\deg(g(x, y))$ with respect to " \leq ". Consequently $r(x, y) = 0$ and so we are done.

(2) Let $0 \neq f(x, y) \in \mathcal{C} = \langle g(x, y) \rangle_l$. Then there exists $q(x, y) \in \mathbb{F}_q[x, y; \rho, \theta]$ such that $f(x, y) = q(x, y) \star g(x, y)$. Notice that $\deg(f(x, y)) \leq (l-1, s-1)$. Hence

$$\begin{aligned} \deg(q(x, y)) &= \deg(f(x, y)) - \deg(g(x, y)) \\ &\leq (l-1, s-1) - (l-k, s-t) = (k-1, t-1). \end{aligned}$$

Therefore, \mathbb{B} is a generating set for \mathcal{C} . Assume that $\{a_{i,j} \mid 0 \leq i \leq k-1, 0 \leq j \leq t-1\}$ is a subset of R such that

$$\sum_{j=0}^{t-1} \sum_{i=0}^{k-1} a_{i,j} x^i y^j \star g(x, y) = 0.$$

Given $r(x, y) := \sum_{j=0}^{t-1} \sum_{i=0}^{k-1} a_{i,j} x^i y^j$ such that $r(x, y) \neq 0$, we have that $\deg(r(x, y)) \leq (k-1, t-1)$ and $r(x, y) \star g(x, y) = 0$. Therefore, there exist $q_1(x, y)$ and $q_2(x, y)$ in $\mathbb{F}_q[x, y; \rho, \theta]$ such that

$$r(x, y) \star g(x, y) = q_1(x, y) \star (x^l - \lambda_1) + q_2(x, y) \star (y^s - \lambda_2).$$

Hence

$$\begin{aligned} \deg(q_1(x, y) \star (x^l - \lambda_1) + q_2(x, y) \star (y^s - \lambda_2)) &= \deg(r(x, y)) + \deg(g(x, y)) \\ &\leq (l-1, s-1), \end{aligned}$$

which is a contradiction. Consequently $r(x, y) = 0$ that implies $a_{i,j} = 0$ for all $0 \leq i \leq k-1$ and $0 \leq j \leq t-1$. \square

Proposition 3.7. *Let R be a domain and λ_1, λ_2 be units in R . Suppose that \mathcal{C} is a 2-D skew (λ_1, λ_2) -constacyclic code in R° that is generated by*

$$g(x, y) = \sum_{j=0}^{s-t-1} \sum_{i=0}^{l-k-1} g_{i,j} x^i y^j + \sum_{j=0}^{s-t} x^{l-k} y^j + \sum_{i=0}^{l-k-1} x^i y^{s-t}.$$

Then $g(x, y) \star xy \in \mathcal{C}$ if and only if $g(x, y) \in R^{\rho\theta}[x, y; \rho, \theta]$.

Proof. Clearly $xy \star g(x, y) \in \mathcal{C}$. Assume that $g(x, y) \star xy \in \mathcal{C}$. Therefore

$$\begin{aligned} xy \star g(x, y) - g(x, y)xy &= \sum_{j=0}^{s-t-1} \sum_{i=0}^{l-k-1} (\rho\theta(g_{i,j}) - g_{i,j}) x^{i+1} y^{j+1} \\ &= p(x, y) \star g(x, y), \end{aligned}$$

for some $p(x, y) \in R[x, y; \rho, \theta]$. It is easy to see that $p(x, y)$ is constant and $p(x, y)g_{0,0} = 0$. Since $g(x, y)$ is a right divisor of $(x^l - \lambda_1) \star (y^s - \lambda_2)$

and λ_1, λ_2 are units, then $g_{0,0}$ is a unit. Hence $p(x, y) = 0$ and so $\rho^\theta(g_{i,j}) = g_{i,j}$ for all $0 \leq i \leq l-k-1$ and $0 \leq j \leq s-t-1$. Therefore $g(x, y) \in R^{\rho^\theta}[x, y; \rho, \theta]$.

For the converse, assume that $g(x, y) \in R^{\rho^\theta}[x, y; \rho, \theta]$. Thus $xy \star g_{i,j} = \rho^\theta(g_{i,j})xy = g_{i,j}xy$ for all $0 \leq i \leq l-k-1$ and $0 \leq j \leq s-t-1$. Hence $g(x, y)xy = xy \star g(x, y) \in \mathcal{C}$, so we are done. \square

Definition 3.8. For two $l \times s$ arrays $c = (c_{i,j})$ and $d = (d_{i,j})$ we define:

$$c \odot d := \sum_{j=0}^{s-1} \sum_{i=0}^{l-1} c_{i,j} d_{i,j}.$$

We say that c, d are orthogonal if $c \odot d = 0$. The dual code of a linear code \mathcal{C} of length ls is the set of all $l \times s$ arrays orthogonal to all codewords of \mathcal{C} . The dual code of \mathcal{C} is denoted by \mathcal{C}^\perp .

Theorem 3.9. Let $|\langle \rho \rangle| \mid l$, $|\langle \theta \rangle| \mid s$ and λ_1, λ_2 be units in R such that $\lambda_1 \in R^\rho, \lambda_2 \in R^\theta$. Suppose that \mathcal{C} is a code of length ls over R . Then \mathcal{C} is 2-D skew (λ_1, λ_2) -constacyclic if and only if \mathcal{C}^\perp is 2-D skew $(\lambda_1^{-1}, \lambda_2^{-1})$ -constacyclic. In particular, if $\lambda_1^2 = \lambda_2^2 = 1$, then \mathcal{C} is 2-D skew (λ_1, λ_2) -constacyclic if and only if \mathcal{C}^\perp is 2-D skew (λ_1, λ_2) -constacyclic.

Proof. First of all, $\lambda_1 \in R^\rho, \lambda_2 \in R^\theta$ imply that $\lambda_1 \in R^{\rho^{-1}}, \lambda_2 \in R^{\theta^{-1}}$. Let $c = (c_{i,j}) \in \mathcal{C}$ and $b = (b_{i,j}) \in \mathcal{C}^\perp$ be two $l \times s$ arrays. Since \mathcal{C} is a column λ_1 -constacyclic code, then

$$\begin{pmatrix} \rho^{l-1}(\lambda_1 c_{1,0}) & \rho^{l-1}(\lambda_1 c_{1,1}) & \cdots & \rho^{l-1}(\lambda_1 c_{1,s-1}) \\ \rho^{l-1}(\lambda_1 c_{2,0}) & \rho^{l-1}(\lambda_1 c_{2,1}) & \cdots & \rho^{l-1}(\lambda_1 c_{2,s-1}) \\ \vdots & \vdots & \ddots & \vdots \\ \rho^{l-1}(\lambda_1 c_{l-1,0}) & \rho^{l-1}(\lambda_1 c_{l-1,1}) & \cdots & \rho^{l-1}(\lambda_1 c_{l-1,s-1}) \\ \rho^{l-1}(c_{0,0}) & \rho^{l-1}(c_{0,1}) & \cdots & \rho^{l-1}(c_{0,s-1}) \end{pmatrix}$$

is in \mathcal{C} and so it is orthogonal to $b = (b_{i,j})$, i.e.,

$$\begin{aligned} 0 &= \sum_{i=1}^{l-1} \sum_{j=0}^{s-1} \rho^{l-1}(\lambda_1 c_{i,j}) b_{i-1,j} + \sum_{j=0}^{s-1} \rho^{l-1}(c_{0,j}) b_{l-1,j} \\ &= \lambda_1 \left(\sum_{i=1}^{l-1} \sum_{j=0}^{s-1} \rho^{l-1}(c_{i,j}) b_{i-1,j} + \sum_{j=0}^{s-1} \rho^{l-1}(c_{0,j}) \lambda_1^{-1} b_{l-1,j} \right) \end{aligned}$$

Therefore

$$\sum_{i=1}^{l-1} \sum_{j=0}^{s-1} \rho^{l-1}(c_{i,j}) b_{i-1,j} + \sum_{j=0}^{s-1} \rho^{l-1}(c_{0,j}) \lambda_1^{-1} b_{l-1,j} = 0.$$

Hence

$$0 = \rho(0) = \sum_{i=1}^{l-1} \sum_{j=0}^{s-1} c_{i,j} \rho(b_{i-1,j}) + \sum_{j=0}^{s-1} c_{0,j} \lambda_1^{-1} \rho(b_{l-1,j}),$$

which shows that

$$\begin{pmatrix} \lambda_1^{-1} \rho(b_{l-1,0}) & \lambda_1^{-1} \rho(b_{l-1,1}) & \cdots & \lambda_1^{-1} \rho(b_{l-1,s-1}) \\ \rho(b_{0,0}) & \rho(b_{0,1}) & \cdots & \rho(b_{0,s-1}) \\ \vdots & \vdots & \ddots & \vdots \\ \rho(b_{l-2,0}) & \rho(b_{l-2,1}) & \cdots & \rho(b_{l-2,s-1}) \end{pmatrix}$$

is orthogonal to $c = (c_{i,j})$ and so it belongs to \mathcal{C}^\perp . Thus \mathcal{C}^\perp is column λ_1^{-1} -constacyclic. On the other hand, since \mathcal{C} is a row λ_2 -constacyclic code, then

$$\begin{pmatrix} \theta^{s-1}(\lambda_2 c_{0,1}) & \cdots & \theta^{s-1}(\lambda_2 c_{0,s-1}) & \theta^{s-1}(c_{0,0}) \\ \theta^{s-1}(\lambda_2 c_{1,1}) & \cdots & \theta^{s-1}(\lambda_2 c_{1,s-1}) & \theta^{s-1}(c_{1,0}) \\ \vdots & \ddots & \vdots & \vdots \\ \theta^{s-1}(\lambda_2 c_{l-1,1}) & \cdots & \theta^{s-1}(\lambda_2 c_{l-1,s-1}) & \theta^{s-1}(c_{l-1,0}) \end{pmatrix}$$

is in \mathcal{C} and so it is orthogonal to $b = (b_{i,j})$, i.e.,

$$\begin{aligned} 0 &= \sum_{j=1}^{s-1} \sum_{i=0}^{l-1} \theta^{s-1}(\lambda_2 c_{i,j}) b_{i,j-1} + \sum_{i=0}^{l-1} \theta^{s-1}(c_{i,0}) b_{i,s-1} \\ &= \lambda_2 \left(\sum_{j=1}^{s-1} \sum_{i=0}^{l-1} \theta^{s-1}(c_{i,j}) b_{i,j-1} + \sum_{i=0}^{l-1} \theta^{s-1}(c_{i,0}) \lambda_2^{-1} b_{i,s-1} \right) \end{aligned}$$

Thus

$$\sum_{j=1}^{s-1} \sum_{i=0}^{l-1} \theta^{s-1}(c_{i,j}) b_{i,j-1} + \sum_{i=0}^{l-1} \theta^{s-1}(c_{i,0}) \lambda_2^{-1} b_{i,s-1} = 0.$$

So we have

$$0 = \theta(0) = \sum_{j=1}^{s-1} \sum_{i=0}^{l-1} c_{i,j} \theta(b_{i,j-1}) + \sum_{i=0}^{l-1} c_{i,0} \lambda_2^{-1} \theta(b_{i,s-1}),$$

which shows that

$$\begin{pmatrix} \lambda_2^{-1} \theta(b_{0,s-1}) & \theta(b_{0,0}) & \cdots & \theta(b_{0,s-2}) \\ \lambda_2^{-1} \theta(b_{1,s-1}) & \theta(b_{1,0}) & \cdots & \theta(b_{1,s-2}) \\ \vdots & \vdots & \ddots & \vdots \\ \lambda_2^{-1} \theta(b_{l-1,s-1}) & \theta(b_{l-1,0}) & \cdots & \theta(b_{l-1,s-2}) \end{pmatrix}$$

is orthogonal to $c = (c_{i,j})$ and so it belongs to \mathcal{C}^\perp . Thus, it follows that \mathcal{C}^\perp is row λ_2^{-1} -constacyclic. Consequently \mathcal{C}^\perp is 2- D skew $(\lambda_1^{-1}, \lambda_2^{-1})$ -constacyclic.

The converse holds by the fact that $(\mathcal{C}^\perp)^\perp = \mathcal{C}$. \square

The ring $R[x, y; \rho, \theta]$ can be localized to the right at the multiplicative set $S = \{x^i y^j \mid i, j \in \mathbb{N}\}$. The existence of the localization $R[x, y; \rho, \theta]S^{-1}$ follows from [19, Theorem 2] since S verifies the following two necessary and sufficient conditions:

- (1) For all $x^{i_1} y^{j_1} \in S$ and $f(x, y) \in R[x, y; \rho, \theta]$, there exists $x^{i_2} y^{j_2} \in S$ and $g(x, y) \in R[x, y; \rho, \theta]$ such that $f(x, y)x^{i_1} y^{j_1} = x^{i_2} y^{j_2} \star g(x, y)$. To prove this note to the multiplication rule $x^i y^j \star a = \rho^i \theta^j (a)x^i y^j$.
- (2) If for $x^{i_1} y^{j_1} \in S$ and $f(x, y) \in R[x, y; \rho, \theta]$ we have $x^{i_1} y^{j_1} \star f(x, y) = 0$, then there exists $x^{i_2} y^{j_2} \in S$ such that $f(x, y)x^{i_2} y^{j_2} = 0$. But since $x^{i_2} y^{j_2}$ is never a zero divisor, $f(x, y)$ must be zero.

Now, we consider the ring $R[x, y; \rho, \theta]S^{-1}$ consisting of the elements $\sum_{j=0}^t \sum_{i=0}^k x^{-i} y^{-j} a_{i,j}$, where the coefficients are on the right and where the multiplication rule is given by $ax^{-1} y^{-1} = x^{-1} y^{-1} \rho \theta (a)$.

Proposition 3.10. *Let $\psi : R[x, y; \rho, \theta] \rightarrow R[x, y; \rho, \theta]S^{-1}$ be defined by*

$$\psi\left(\sum_{j=0}^t \sum_{i=0}^k a_{i,j} x^i y^j\right) = \sum_{j=0}^t \sum_{i=0}^k x^{-i} y^{-j} a_{i,j}.$$

Then ψ is a ring anti-isomorphism.

Proof. Similar to [4, Theorem 4.4]. \square

From now on, we assume that $\lambda_1, \lambda_2 \in R^{\rho, \theta}$, $|\langle \rho \rangle| = l$ and $|\langle \theta \rangle| = s$.

Theorem 3.11. *Let $a(x, y) = \sum_{j=0}^{s-1} \sum_{i=0}^{l-1} a_{i,j} x^i y^j$ and $b(x, y) = \sum_{j=0}^{s-1} \sum_{i=0}^{l-1} b_{i,j} x^i y^j$ be in $R[x, y; \rho, \theta]$. Suppose that $\lambda_1^2 = \lambda_2^2 = 1$. Then the following conditions are equivalent:*

- (1) *The coefficient matrix of $a(x, y)$ is orthogonal to the coefficient matrix of $x^i y^j (x^{l-1} y^{s-1} \star \psi(b(x, y)))$ for all $i \in \{0, 1, \dots, l-1\}$ and all $j \in \{0, 1, \dots, s-1\}$;*

(2) The coefficient matrix of $a(x, y)$ is orthogonal to

$$\mathcal{A} = \begin{pmatrix} b_{l-1,s-1} & \theta(b_{l-1,s-2}) & \cdots & \theta^{s-1}(b_{l-1,0}) \\ \rho(b_{l-2,s-1}) & \rho\theta(b_{l-2,s-2}) & \cdots & \rho\theta^{s-1}(b_{l-2,0}) \\ \vdots & \vdots & \ddots & \vdots \\ \rho^{l-1}(b_{0,s-1}) & \rho^{l-1}\theta(b_{0,s-2}) & \cdots & \rho^{l-1}\theta^{s-1}(b_{0,0}) \end{pmatrix}$$

and all of its column skew λ_1 -constacyclic shifts and row skew λ_2 -constacyclic shifts;

(3) $a(x, y) \star b(x, y) = 0$ in R° .

Proof. (1) \Leftrightarrow (2) With a routine computation we can deduce that the coefficient matrix of $x^{l-1}y^{s-1} \star \psi(b(x, y))$ is equal to \mathcal{A} , also the coefficient matrices of $x^i y^j (x^{l-1}y^{s-1} \star \psi(b(x, y)))$ for $0 \leq i \leq l-1$ and $0 \leq j \leq s-1$ are precisely column skew λ_1 -constacyclic shifts and row skew λ_2 -constacyclic shifts of \mathcal{A} .

(2) \Leftrightarrow (3) Suppose that $a(x, y) \star b(x, y) = \sum_{t=0}^{s-1} \sum_{k=0}^{l-1} c_{i,j} x^i y^j \in R^\circ$. For $k \in \{0, 1, \dots, l-1\}$ and $t \in \{0, 1, \dots, s-1\}$ we have that

$$\begin{aligned} c_{k,t} &= \sum_{\substack{j+v=t \\ 0 \leq j, v \leq s-1}} \sum_{\substack{i+u=k \\ 0 \leq i, u \leq l-1}} a_{i,j} \rho^i \theta^j (b_{uv}) + \sum_{\substack{j+v=t \\ 0 \leq j, v \leq s-1}} \sum_{\substack{i+u=k+l \\ 0 \leq i, u \leq l-1}} a_{i,j} \rho^i \theta^j (\lambda_1 b_{uv}) \\ &+ \sum_{\substack{j+v=t+s \\ 0 \leq j, v \leq s-1}} \sum_{\substack{i+u=k \\ 0 \leq i, u \leq l-1}} a_{i,j} \rho^i \theta^j (\lambda_2 b_{uv}) + \sum_{\substack{j+v=t+s \\ 0 \leq j, v \leq s-1}} \sum_{\substack{i+u=k+l \\ 0 \leq i, u \leq l-1}} a_{i,j} \rho^i \theta^j (\lambda_1 \lambda_2 b_{uv}) \\ &= \lambda_1 \lambda_2 \left(\sum_{\substack{j+v=t \\ 0 \leq j, v \leq s-1}} \sum_{\substack{i+u=k \\ 0 \leq i, u \leq l-1}} a_{i,j} \rho^{k-u} \theta^{t-v} (\lambda_1 \lambda_2 b_{uv}) \right. \\ &+ \sum_{\substack{j+v=t \\ 0 \leq j, v \leq s-1}} \sum_{\substack{i+u=k+l \\ 0 \leq i, u \leq l-1}} a_{i,j} \rho^{k+l-u} \theta^{t-v} (\lambda_2 b_{uv}) \\ &+ \sum_{\substack{j+v=t+s \\ 0 \leq j, v \leq s-1}} \sum_{\substack{i+u=k \\ 0 \leq i, u \leq l-1}} a_{i,j} \rho^{k-u} \theta^{t+s-v} (\lambda_1 b_{uv}) \\ &+ \left. \sum_{\substack{j+v=t+s \\ 0 \leq j, v \leq s-1}} \sum_{\substack{i+u=k+l \\ 0 \leq i, u \leq l-1}} a_{i,j} \rho^{k+l-u} \theta^{t+s-v} (b_{uv}) \right) \end{aligned}$$

which is the \odot -product of

$$\lambda_1 \lambda_2 \begin{pmatrix} a_{0,0} & \cdots & a_{0,s-1} \\ a_{1,0} & \cdots & a_{1,s-1} \\ \vdots & \ddots & \vdots \\ a_{l-1,0} & \cdots & a_{l-1,s-1} \end{pmatrix}$$

in the matrix

$$\begin{pmatrix} \lambda_1 \lambda_2 b_{k,t} & \theta(\lambda_1 \lambda_2 b_{k,t-1}) \dots \theta^t(\lambda_1 \lambda_2 b_{k,0}) & \theta^{t+1}(\lambda_1 b_{k,s-1}) \dots \theta^{s-1}(\lambda_1 b_{k,t+1}) \\ \rho(\lambda_1 \lambda_2 b_{k-1,t}) & \rho\theta(\lambda_1 \lambda_2 b_{k-1,t-1}) \dots \rho\theta^t(\lambda_1 \lambda_2 b_{k-1,0}) & \rho\theta^{t+1}(\lambda_1 b_{k-1,s-1}) \dots \rho\theta^{s-1}(\lambda_1 b_{k-1,t+1}) \\ \vdots & \vdots & \vdots \\ \rho^k(\lambda_1 \lambda_2 b_{0,t}) & \rho^k\theta(\lambda_1 \lambda_2 b_{0,t-1}) \dots \rho^k\theta^t(\lambda_1 \lambda_2 b_{0,0}) & \rho^k\theta^{t+1}(\lambda_1 b_{0,s-1}) \dots \rho^k\theta^{s-1}(\lambda_1 b_{0,t+1}) \\ \rho^{k+1}(\lambda_2 b_{l-1,t}) & \rho^{k+1}\theta(\lambda_2 b_{l-1,t-1}) \dots \rho^{k+1}\theta^t(\lambda_2 b_{l-1,0}) & \rho^{k+1}\theta^{t+1}(b_{l-1,s-1}) \dots \rho^{k+1}\theta^{s-1}(b_{l-1,t+1}) \\ \vdots & \vdots & \vdots \\ \rho^{l-1}(\lambda_2 b_{k+1,t}) & \rho^{l-1}\theta(\lambda_2 b_{k+1,t-1}) \dots \rho^{l-1}\theta^t(\lambda_2 b_{k+1,0}) & \rho^{l-1}\theta^{t+1}(b_{k+1,s-1}) \dots \rho^{l-1}\theta^{s-1}(b_{k+1,t+1}) \end{pmatrix}.$$

Therefore $a(x, y) \star b(x, y) = 0$ in R° if and only if $c_{k,t} = 0$ for all $k \in \{0, 1, \dots, l-1\}$ and all $t \in \{0, 1, \dots, s-1\}$ which is true if and only if

$$\begin{pmatrix} a_{0,0} & \dots & a_{0,s-1} \\ a_{1,0} & \dots & a_{1,s-1} \\ \vdots & \ddots & \vdots \\ a_{l-1,0} & \dots & a_{l-1,s-1} \end{pmatrix}$$

is orthogonal to \mathcal{A} and all of its column skew λ_1 -constacyclic shifts and row skew λ_2 -constacyclic shifts. \square

Proposition 3.12. *Let $\lambda_1^2 = \lambda_2^2 = 1$, $g(x, y)$ be a right divisor of $(x^l - \lambda_1) \star (y^s - \lambda_2)$ in $R[x, y; \rho, \theta]$ and $h(x, y) := \frac{(x^l - \lambda_1) \star (y^s - \lambda_2)}{g(x, y)}$ with $\deg(h(x, y)) = (k, t)$. Suppose that \mathcal{C} is a 2-D skew (λ_1, λ_2) -constacyclic code of length ls over R that is generated by $g(x, y)$. Then the bivariate skew polynomial $x^k y^t \star \psi(h(x, y))$ is a right divisor of $(x^l - \lambda_1) \star (y^s - \lambda_2)$ and $x^k y^t \star \psi(h(x, y)) \in \mathcal{C}^\perp$.*

Proof. By the assumptions that $\lambda_1, \lambda_2 \in R^{\rho, \theta}$, $|\langle \rho \rangle| \mid l$ and $|\langle \theta \rangle| \mid s$ we can see that

$$\begin{aligned} (\psi(g(x, y)) \star \lambda_1 \lambda_2 x^{l-k} y^{s-t}) \star (x^k y^t \star \psi(h(x, y))) &= \psi(g(x, y)) \star \lambda_1 \lambda_2 x^l y^s \star \psi(h(x, y)) \\ &= \lambda_1 \lambda_2 x^l y^s \star \psi(g(x, y)) \star \psi(h(x, y)) \\ &= \lambda_1 \lambda_2 x^l y^s \star \psi(h(x, y) \star g(x, y)) \\ &\text{(since } \psi \text{ is a ring anti-isomorphism)} \\ &= \lambda_1 \lambda_2 x^l y^s \star \psi((x^l - \lambda_1) \star (y^s - \lambda_2)) \\ &= \lambda_1 \lambda_2 x^l y^s \star (x^{-l} y^{-s} \\ &\quad - x^{-l} \lambda_2 - y^{-s} \lambda_1 + \lambda_1 \lambda_2) \\ &= (x^l - \lambda_1) \star (y^s - \lambda_2). \end{aligned}$$

Moreover, since $g(x, y) \star h(x, y) = 0$ in R° , then Theorem 3.11 implies that $x^k y^t \star \psi(h(x, y)) \in \mathcal{C}^\perp$. \square

Now we can state the following open problem.

Problem 3.13. *In Proposition 3.12, is \mathcal{C}^\perp generated by $x^k y^t \star \psi(h(x, y))$?*

Theorem 3.14. *Let $g(x, y)$ be a monic right divisor of $(x^l - \lambda_1) \star (y^s - \lambda_2)$ in $R[x, y; \rho, \theta]$ and $h(x, y) := \frac{(x^l - \lambda_1) \star (y^s - \lambda_2)}{g(x, y)}$. Suppose that \mathcal{C} is a 2- D skew (λ_1, λ_2) -constacyclic code of length ls over R that is generated by $g(x, y)$. Then for $f(x, y) \in R[x, y; \rho, \theta]$, $f(x, y) \in \mathcal{C}$ if and only if $f(x, y) \star h(x, y) = 0$ in $R^\diamond = R[x, y; \rho, \theta] / \langle (x^l - \lambda_1) \star (y^s - \lambda_2) \rangle_l$.*

Proof. Let $f(x, y) \in \mathcal{C}$. Then $f(x, y) = q(x, y) \star g(x, y)$ for some $q(x, y) \in R[x, y; \rho, \theta]$. So we have

$$\begin{aligned} f(x, y) \star h(x, y) &= q(x, y) \star g(x, y) \star h(x, y) \\ &= q(x, y) \star (x^l - \lambda_1) \star (y^s - \lambda_2) = 0, \end{aligned}$$

in R^\diamond . Conversely, assume that $f(x, y) \star h(x, y) = 0$ in R^\diamond . Then $f(x, y) \star h(x, y) = q(x, y) \star (x^l - \lambda_1) \star (y^s - \lambda_2)$ for some $q(x, y) \in R[x, y; \rho, \theta]$. Therefore $f(x, y) \star h(x, y) = q(x, y) \star g(x, y) \star h(x, y)$. Now, since $h(x, y)$ is monic, $f(x, y) = q(x, y) \star g(x, y) \in \mathcal{C}$. \square

Acknowledgments

The author would like to thank the referee for constructive comments which will help to improve the quality of the paper.

REFERENCES

1. M.C. V. Amarra and F.R. Nemenzo, *On $(1 - u)$ -cyclic codes over $\mathbb{F}_{p^k} + u\mathbb{F}_{p^k}$* , App. Math. Letters, **21** (2008), 1129–1133.
2. A. Bonnecaze and P. Udaya, *Cyclic codes and self-dual codes over $\mathbb{F}_2 + u\mathbb{F}_2$* , IEEE Trans. Inf. Theory, **45** (1999), 1250–1255.
3. D. Boucher, W. Geiselmann and F. Ulmer, *Skew-cyclic codes*, Appl. Algebra Eng., Commun. Comput, **18** (4) (2007), 379–389.
4. D. Boucher, P. Sole and F. Ulmer, *Skew constacyclic codes over Galois rings*, Adv. Math. Commun, **2** (3) (2008), 273–292.
5. D. Boucher and F. Ulmer, *Codes as modules over skew polynomial rings*, Lect. Notes Comp. Sci, **5921** (2009), 38–55.
6. D. Boucher and F. Ulmer, *Coding with skew polynomial rings*, J. Symb. Comput, **44** (2009), 1644–1656.
7. H.Q. Dinh, *Constacyclic codes of length 2^s over Galois extension rings of $\mathbb{F}_2 + u\mathbb{F}_2$* , IEEE Trans. Inf. Theory, **55** (2009), 1730–1740.
8. H.Q. Dinh, *Negacyclic codes of length 2^s over Galois rings*, IEEE Trans. Inf. Theory, **51** (2005), 4252–4262.
9. H.Q. Dinh, *Constacyclic codes of length p^s over $\mathbb{F}_{p^m} + u\mathbb{F}_{p^m}$* , J. Algebra, **324** (2010), 940–950.
10. H.Q. Dinh and S.R. López-Permouth, *Cyclic and negacyclic codes over finite chain rings*, IEEE Trans. Inf. Theory, **50** (2004), 1728–1744.

11. C. Güneri and F. Özbudak, *A relation between quasi-cyclic codes and 2-D cyclic codes*, Finite Fields Appl, **18** (2012), 123–132.
12. A. Hammons, P.V. Kumar, A.R. Calderbank, N.J. A. Sloane and P. Sole, *The \mathbb{Z}_4 linearity of Kerdock, Preparata, Goethals and related codes*, IEEE Trans. Inf. Theory, **40** (4) (1994), 301–319.
13. T. Ikai, H. Kosako and Y. Kojima, *Two-dimensional cyclic codes*, Electron. Commun. Jpn, **57A** (1975), 27–35.
14. H. Imai, *A theory of two-dimensional cyclic codes*, Inf. Control, **34** (1977), 1–21.
15. S. Jitman, S. Ling and P. Udomkavanich, *Skew constacyclic codes over finite chain rings*, Adv. Math. Commun, **6**(1) (2012), 39–63.
16. N.H. McCoy, *Annihilators in polynomial rings*, Amer. Math. Monthly, **64** (1957), 28–29.
17. F.J. MacWilliams and N. J. A. Sloane, *The theory of error correcting codes*, North Holland, 1977.
18. J.F. Qian, L.N. Zhang and S.X. Zhu, *$(1 + u)$ -cyclic and cyclic codes over the ring $\mathbb{F}_2 + u\mathbb{F}_2$* , Appl. Math. Letters, **19** (2006), 820–823.
19. P. Ribenboim., *Sur la localisation des anneaux non commutatifs*, Séminaire Dubreil. Algèbre et théorie des nombres, tome 24. (1970–1971).
20. K. Saints, and C. Heegard, *Algebraic-geometric codes and multidimensional cyclic codes: a unified theory and algorithms for decoding using Gröbner bases*, IEEE Trans. Inf. Theory IT-41, **5** (1993), 1733–1751.
21. S. Sakata, *Decoding binary 2-D cyclic codes by the 2-D Berlekamp-Massey algorithm*, IEEE Trans. Inf. Theory IT-21, **37** (1991), 1200–1203.
22. S. Sakata, *General theory of doubly periodic arrays over an arbitrary finite field and its applications*, IEEE Trans. Inf. Theory IT-24, **6** (1978), 719–730.
23. S. Sakata, *On determining the independent point set for doubly periodic arrays and encoding two-dimensional cyclic codes and their duals*, IEEE Trans. Inf. Theory IT-21, **5** (1981), 556–565.
24. P. Udaya and A. Bonnezeze, *Decoding of cyclic codes over $\mathbb{F}_2 + u\mathbb{F}_2$* , IEEE Trans. Inf. Theory, **45** (1999), 2148–2157.
25. L. Xiuli and L. Hongyan, *2-D skew-cyclic codes over $\mathbb{F}_q[x, y; \rho, \theta]$* , Finite Fields Appl, **25** (2014), 49–63.

Hojjat Mostafanasab

Eski Silahtarağa Elektrik Santrali, Kazim Karabekir, Istanbul Bilgi University, Cad.

No: 2/1334060, Eyüp Istanbul, Turkey.

Email: h.mostafanasab@gmail.com