

ON THE m -EXTENSION DUAL COMPLEX FIBONACCI p -NUMBERS

B. PRASAD *

ABSTRACT. In this paper, we introduced m -extension dual complex Fibonacci p -numbers. We established the properties of m -extension dual complex Fibonacci p -numbers. They are connected to complex Fibonacci numbers, complex Fibonacci p -numbers and dual complex Fibonacci p -numbers.

1. INTRODUCTION

The Fibonacci numbers, F_n are defined by the recurrence relation:

$$F_n = F_{n-1} + F_{n-2} \text{ for } n > 2$$

with initial terms

$$F_1 = F_2 = 1.$$

The Fibonacci numbers, F_n and golden mean (golden ratio),

$$\tau = \lim_{n \rightarrow \infty} \frac{F_n}{F_{n-1}} = \frac{1 + \sqrt{5}}{2}$$

have materialized in several stream of arts, sciences, computer sciences, high energy physics, information and coding theory [4, 6, 7, 15, 16].

MSC(2010): Primary: 11B39; Secondary: 94B35, 94B25.

Keywords: Dual complex Fibonacci p -numbers, Complex Fibonacci p -numbers, Fibonacci numbers, Golden mean.

Received: 29 July 2021, Accepted: 09 September 2021.

*Corresponding author .

The Fibonacci p -numbers [15] are defined by the recurrence relation:

$$F_p(n) = F_p(n-1) + F_p(n-p-1)$$

with $n > p+1$, for a given integer $p = 0, 1, 2, 3, \dots$ and initial terms

$$F_p(1) = F_p(2) = \dots = F_p(p) = F_p(p+1) = 1.$$

The Fibonacci p -numbers, $F_p(n)$ coincide with classical Fibonacci numbers, F_n for $p = 1$. e.g. $F_1(n) = F_n$.

E. Kocer EG et al., [9] introduced the m -extension of Fibonacci p -numbers which satisfy the recurrence relation

$$F_{p,m}(n) = mF_{p,m}(n-1) + F_{p,m}(n-p-1)$$

with initial terms

$$F_{p,m}(1) = a_1, F_{p,m}(2) = a_2, F_{p,m}(3) = a_3, \dots, F_{p,m}(p+1) = a_{p+1}$$

where $p(\geq 0)$ is integer, $m(> 0)$, $n > p+1$ and $a_1, a_2, a_3, \dots, a_{p+1}$ are arbitrary real or complex numbers.

A complex number is of the form $a + ib$ where a and b are real numbers and $i^2 = -1$. It is usually represented by z . The value a is called the real part which is denoted by $Re(z)$ and b is called the imaginary part and denoted by $Im(z)$.

The complex Fibonacci numbers [12] are defined by the recurrence relation:

$$F_n^* = F_{n-1}^* + F_{n-2}^* \text{ for } n \geq 2$$

with initial terms

$$F_0^* = i, F_1^* = 1 + i$$

where i is the imaginary unit which satisfies $i^2 = -1$ and

$$F_n^* = F_n + iF_{n+1}.$$

Clifford Algebra is a powerful mathematical tool that offers a natural and direct way to model geometric objects and their transformations. It makes geometric objects (points, lines and planes) into basic elements of computation and defines few universal operators that are applicable to all types of geometric elements. In 19th century, Clifford [5] defined new number system by using the notion $\varepsilon^2 = 0$ and $\varepsilon \neq 0$ i.e. ε is an nilpotent number. This number system is known as dual number system and dual number are represented in the form $A = a + \varepsilon a^*$ where a and a^* are real numbers and ε is an nilpotent number. Kotelnikov [10] and Study [17] generalized the first applications of dual numbers in mechanics. It has also applications in algebraic geometry, kinematics,

number theory, theory of relativity etc. [11, 2, 13].
Dual complex numbers [11] are defined as follows:

$$W = \{w = z_1 + \varepsilon z_2 : z_1, z_2 \in \mathbf{C} \text{ where } \varepsilon^2 = 0 \text{ and } \varepsilon \neq 0 \}.$$

If $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$ then any dual complex number can be written as follows:

$$w = x_1 + iy_1 + \varepsilon x_2 + i\varepsilon y_2 \text{ where } i^2 = -1, \varepsilon \neq 0, \varepsilon^2 = 0, (i\varepsilon)^2 = 0.$$

The real and dual quaternions form a division ring but dual complex numbers form a commutative ring with characteristics zero. The multiplication of dual complex numbers gives the structure of two dimensional complex Clifford Algebra and four dimensional real Clifford Algebra. The base elements of dual complex numbers satisfies the commutative multiplication scheme which are shown in the **Table 1**.

Table 1: Multiplication scheme of dual complex numbers

x	1	i	ε	$i\varepsilon$
1	1	i	ε	$i\varepsilon$
i	i	-1	$i\varepsilon$	$-\varepsilon$
ε	ε	$i\varepsilon$	0	0
$i\varepsilon$	$i\varepsilon$	$-\varepsilon$	0	0

Five different conjugation operates on dual complex numbers [11] in the following manner:

$$\text{Let } w = z_1 + \varepsilon z_2 = x_1 + iy_1 + \varepsilon x_2 + i\varepsilon y_2, z_2 \neq 0,$$

$$w^{\star 1} = (x_1 - iy_1) + \varepsilon(x_2 - iy_2) = z_1^* + \varepsilon z_2^*, \text{ Complex-conjugation,}$$

$$w^{\star 2} = (x_1 + iy_1) - \varepsilon(x_2 + iy_2) = z_1 - \varepsilon z_2, \text{ Dual-conjugation,}$$

$$w^{\star 3} = (x_1 - iy_1) - \varepsilon(x_2 - iy_2) = z_1^* - \varepsilon z_2^*, \text{ Coupled-conjugation,}$$

$$w^{\star 4} = (x_1 - iy_1)\left(1 - \frac{\varepsilon(x_2 + iy_2)}{(x_1 + iy_1)}\right) = z_1^*\left(1 - \frac{\varepsilon z_2}{z_1}\right), \text{ Dual-complex-conjugation,}$$

$$w^{\star 5} = (x_2 + iy_2) - \varepsilon(x_1 + iy_1) = z_2 - \varepsilon z_1, \text{ Anti-dual-conjugation.}$$

The norm of dual complex numbers defined in the following manner:

$$N_{w^{\star 1}} = \|w \times w^{\star 1}\| = \sqrt{|z_1^2| + 2\varepsilon \text{Re}(z_1 z_2^*)},$$

$$N_{w^{\star 2}} = \|w \times w^{\star 2}\| = \sqrt{|z_1^2|},$$

$$N_{w^{\star 3}} = \|w \times w^{\star 3}\| = \sqrt{|z_1^2| - 2i\varepsilon \text{Im}(z_1 z_2^*)},$$

$$N_{w^{\star 4}} = \|w \times w^{\star 4}\| = \sqrt{|z_1^2|},$$

$$N_{w^{\star 5}} = \|w \times w^{\star 5}\| = \sqrt{z_1 z_2 + \varepsilon(z_2^2 - z_1^2)}.$$

In 2017, Gungor and Azak [8] defined the dual complex Fibonacci numbers as follow: $DF_n^* = F_n + iF_{n+1} + \varepsilon(F_{n+2} + iF_{n+3})$ where the basis $\{1, i, \varepsilon, i\varepsilon\}$ satisfies the condition

$$i^2 = -1, \varepsilon \neq 0, \varepsilon^2 = 0, (i\varepsilon)^2 = 0.$$

In 2018, Aydin Torunbalci [1] defined the dual complex k -Fibonacci

numbers as follow:

$$DCF_{k,n} = F_{k,n} + iF_{k,n+1} + \varepsilon(F_{k,n+2} + iF_{k,n+3})$$

where the basis $\{1, i, \varepsilon, i\varepsilon\}$ satisfies the condition $i^2 = -1, \varepsilon \neq 0, \varepsilon^2 = 0, (i\varepsilon)^2 = 0$.

In 2019, Prasad [12] introduced complex Fibonacci p -numbers by the following recurrence relation:

$$F_p^*(n) = F_p^*(n-1) + F_p^*(n-p-1)$$

with $n > p+1$ and initial terms

$$F_p^*(0) = i, F_p^*(1) = F_p^*(2) = \dots = F_p^*(p) = 1 + i.$$

In 2021, Prasad [14] introduced dual complex Fibonacci p -numbers by the following recurrence relation:

$$DF_p^*(n) = F_p^*(n-1) + \varepsilon F_p^*(n-p-1) = F_p(n-1) + iF_p(n) + \varepsilon(F_p(n-p-1) + iF_p(n-p))$$

with $n > p+1$ and initial terms

$$F_p^*(0) = i, F_p^*(1) = F_p^*(2) = \dots = F_p^*(p) = 1 + i,$$

and the basis $\{1, i, \varepsilon, i\varepsilon\}$ satisfies the condition

$$i^2 = -1, \varepsilon \neq 0, \varepsilon^2 = 0, (i\varepsilon)^2 = 0$$

In this paper, we introduce m -extension dual complex Fibonacci p -numbers by the following recurrence relation:

$$\begin{aligned} DF_{p,m}^*(n) &= mF_{p,m}^*(n-1) + \varepsilon F_{p,m}^*(n-p-1) \quad (1.1) \\ &= m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p)) \end{aligned}$$

with $n > p+1$ and initial terms

$$F_{p,m}^*(n) = m^{n-1}, n = 1, 2, 3, 4, \dots, p+1$$

and the basis $\{1, i, \varepsilon, i\varepsilon\}$ satisfies the condition

$$i^2 = -1, \varepsilon \neq 0, \varepsilon^2 = 0, (i\varepsilon)^2 = 0.$$

2. ON THE m -EXTENSION DUAL COMPLEX FIBONACCI p -NUMBERS

Definition 2.1. On the m -extension dual complex Fibonacci p -numbers are given by the following recurrence relation:

$$\begin{aligned} DF_{p,m}^*(n) &= mF_{p,m}^*(n-1) + \varepsilon F_{p,m}^*(n-p-1) \\ &= m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p)) \\ &= mF_{p,m}(n-1) + imF_{p,m}(n) + \varepsilon F_{p,m}(n-p-1) + i\varepsilon F_{p,m}(n-p) \quad (2.1) \end{aligned}$$

with $n > p + 1$ and initial terms

$$F_{p,m}^*(n) = m^{n-1}, n = 1, 2, 3, 4, \dots, p + 1$$

and the basis $\{1, i, \varepsilon, i\varepsilon\}$ satisfies the condition $i^2 = -1, \varepsilon \neq 0, \varepsilon^2 = 0, (i\varepsilon)^2 = 0$.

With the addition and multiplication by real scalar of two m -extension dual complex Fibonacci p -numbers gives the m -extension dual complex Fibonacci p -numbers.

Definition 2.2. The addition and subtraction of the m -extension dual complex Fibonacci p -numbers are defined by

$$\begin{aligned} DF_{p,m}^*(n_1) \pm DF_{p,m}^*(n_2) &= mF_{p,m}^*(n_1 - 1) + \varepsilon F_{p,m}^*(n_1 - p - 1) \pm mF_{p,m}^*(n_2 - 1) \\ &+ \varepsilon F_{p,m}^*(n_2 - p - 1) \\ &= m(F_{p,m}(n_1 - 1) + iF_{p,m}(n_1)) + \varepsilon(F_{p,m}(n_1 - p - 1) + iF_{p,m}(n_1 - p)) \pm \\ &m(F_{p,m}(n_2 - 1) + iF_{p,m}(n_2)) + \varepsilon(F_{p,m}(n_2 - p - 1) + iF_{p,m}(n_2 - p)) \\ &= m(F_{p,m}(n_1 - 1) \pm F_{p,m}(n_2 - 1)) + im(F_{p,m}(n_1) \pm F_{p,m}(n_2)) + \varepsilon(F_{p,m}(n_1 - p - 1) \\ &\pm F_{p,m}(n_2 - p - 1)) + i\varepsilon(F_{p,m}(n_1 - p) \pm F_{p,m}(n_2 - p)). \end{aligned}$$

Definition 2.3. The multiplication of a m -extension dual complex Fibonacci p -number by real scalar λ is defined by

$$\begin{aligned} \lambda DF_{p,m}^*(n) &= \lambda mF_{p,m}(n - 1) + i\lambda mF_{p,m}(n) + \varepsilon\lambda F_{p,m}(n - p - 1) + \\ & i\varepsilon\lambda F_{p,m}(n - p). \end{aligned}$$

Definition 2.4. The multiplication of two m -extension dual complex Fibonacci p -numbers is defined by

$$\begin{aligned} DF_{p,m}^*(n_1)DF_{p,m}^*(n_2) &= [mF_{p,m}(n_1 - 1) + imF_{p,m}(n_1) + \varepsilon F_{p,m}(n_1 - p - 1) \\ &+ i\varepsilon F_{p,m}(n_1 - p)][mF_{p,m}(n_2 - 1) + imF_{p,m}(n_2) + \varepsilon F_{p,m}(n_2 - p - 1) + \\ &i\varepsilon F_{p,m}(n_2 - p)] \\ &= m^2[F_{p,m}(n_1 - 1)F_{p,m}(n_2 - 1) - F_{p,m}(n_1)F_{p,m}(n_2)] + im[F_{p,m}(n_1 - 1) \\ &F_{p,m}(n_2) + F_{p,m}(n_1)F_{p,m}(n_2 - 1)] + m\varepsilon[F_{p,m}(n_1 - 1)F_{p,m}(n_2 - p - 1) + \\ &F_{p,m}(n_1 - p - 1)F_{p,m}(n_2 - 1) - F_{p,m}(n_1)F_{p,m}(n_2 - p) - F_{p,m}(n_2)F_{p,m}(n_1 - p)] \\ &+ im\varepsilon[F_{p,m}(n_1 - 1)F_{p,m}(n_2 - p) + F_{p,m}(n_1)F_{p,m}(n_2 - p - 1) + F_{p,m}(n_2)F_{p,m}(n_1 \\ &- p - 1) + F_{p,m}(n_1 - p)F_{p,m}(n_2 - 1)] = DF_{p,m}^*(n_2)DF_{p,m}^*(n_1). \end{aligned}$$

On the m -extension dual complex Fibonacci p -numbers also provide the properties of norm and conjugation.

Proposition 2.5. *Five kinds of conjugation can be defined for dual complex numbers [11]. Therefore, conjugation of the m -extension dual complex Fibonacci p -number is defined in five different ways in the*

following manner:

$$DF_{p,m}^{\star 1}(n) = m(F_{p,m}(n-1) - iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) - iF_{p,m}(n-p)), \text{Complex - conjugation} \quad (2.2)$$

$$DF_{p,m}^{\star 2}(n) = m(F_{p,m}(n-1) + iF_{p,m}(n)) - \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p)), \text{Dual - conjugation} \quad (2.3)$$

$$DF_{p,m}^{\star 3}(n) = m(F_{p,m}(n-1) - iF_{p,m}(n)) - \varepsilon(F_{p,m}(n-p-1) - iF_{p,m}(n-p)), \text{Coupled - conjugation} \quad (2.4)$$

$$DF_{p,m}^{\star 4}(n) = m(F_{p,m}(n-1) - iF_{p,m}(n)) \left(1 - \varepsilon \frac{F_{p,m}(n-p-1) + iF_{p,m}(n-p)}{m(F_{p,m}(n-1) + iF_{p,m}(n))}\right), \text{Dual - complex - conjugation} \quad (2.5)$$

$$DF_{p,m}^{\star 5}(n) = m(F_{p,m}(n-p-1) + iF_{p,m}(n-p)) - \varepsilon(F_{p,m}(n-1) + iF_{p,m}(n)), \text{Anti - dual - conjugation} \quad (2.6)$$

Some properties of conjugations for m -extension dual complex Fibonacci p -numbers are given in the following theorems.

Theorem 2.6. Let $DF_{p,m}^{\star 1}(n)$, $DF_{p,m}^{\star 2}(n)$, $DF_{p,m}^{\star 3}(n)$, $DF_{p,m}^{\star 4}(n)$, $DF_{p,m}^{\star 5}(n)$ be five kinds of conjugation of m -extension dual complex Fibonacci p -number. Thereby, we give the following relations.

$$DF_{p,m}^{\star}(n)DF_{p,m}^{\star 1}(n) = m^2[F_{p,m}^2(n-1) + F_{p,m}^2(n)] + 2m\varepsilon[F_{p,m}(n-1)F_{p,m}(n-p-1) + F_{p,m}(n)F_{p,m}(n-p)] \quad (2.7)$$

$$DF_{p,m}^{\star}(n)DF_{p,m}^{\star 2}(n) = m^2[F_{p,m}^2(n-1) - F_{p,m}^2(n)] + i2m^2F_{p,m}(n-1)F_{p,m}(n) \quad (2.8)$$

$$DF_{p,m}^{\star}(n)DF_{p,m}^{\star 3}(n) = m^2[F_{p,m}^2(n-1) + F_{p,m}^2(n)] + i2m\varepsilon[F_{p,m}(n-p)F_{p,m}(n-1) - F_{p,m}(n)F_{p,m}(n-p-1)] \quad (2.9)$$

$$DF_{p,m}^{\star}(n)DF_{p,m}^{\star 4}(n) = m^2[F_{p,m}^2(n-1) + F_{p,m}^2(n)] \quad (2.10)$$

$$\begin{aligned}
 DF_{p,m}^*(n)DF_{p,m}^{*5}(n) &= m^2[F_{p,m}(n-1)F_{p,m}(n-p-1) - F_{p,m}(n)F_{p,m}(n-p)] \\
 &\quad + im^2[F_{p,m}(n-1)F_{p,m}(n-p) + F_{p,m}(n)F_{p,m}(n-p-1)] \\
 &\quad + m\varepsilon[F_{p,m}^2(n-p-1) + F_{p,m}^2(n) - F_{p,m}^2(n-1) - F_{p,m}^2(n-p)] \\
 &\quad + i2m\varepsilon[F_{p,m}(n-p-1)F_{p,m}(n-p) - F_{p,m}(n-1)F_{p,m}(n)] \quad (2.11)
 \end{aligned}$$

$$DF_{p,m}^*(n) + DF_{p,m}^{*1}(n) = 2[mF_{p,m}(n-1) + \varepsilon F_{p,m}(n-p-1)] \quad (2.12)$$

$$DF_{p,m}^*(n) + DF_{p,m}^{*2}(n) = 2m[F_{p,m}(n-1) + iF_{p,m}(n)] \quad (2.13)$$

$$DF_{p,m}^*(n) + DF_{p,m}^{*3}(n) = 2[mF_{p,m}(n-1) + i\varepsilon F_{p,m}(n-p)] \quad (2.14)$$

$$(F_{p,m}(n-1) + iF_{p,m}(n))DF_{p,m}^{*4}(n) = [F_{p,m}(n-1) - iF_{p,m}(n)]DF_{p,m}^{*2}(n) \quad (2.15)$$

$$\varepsilon DF_{p,m}^*(n) + mDF_{p,m}^{*5}(n) = m^2[F_{p,m}(n-p-1) + iF_{p,m}(n-p)] \quad (2.16)$$

$$mDF_{p,m}^*(n) - \varepsilon DF_{p,m}^{*5}(n) = m^2[F_{p,m}(n-1) + iF_{p,m}(n)] \quad (2.17)$$

$$DF_{p,m}^*(n) + DF_{p,m}^*(n+p) = DF_{p,m}^*(n+p+1) \quad (2.18)$$

Proof. $DF_{p,m}^*(n)DF_{p,m}^{*1}(n) = [m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))][m(F_{p,m}(n-1) - iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) - iF_{p,m}(n-p))] = m^2[F_{p,m}^2(n-1) + F_{p,m}^2(n)] + 2m\varepsilon[F_{p,m}(n-1)F_{p,m}(n-p-1) + F_{p,m}(n)F_{p,m}(n-p)]$, by equations (2) and (3).

$DF_{p,m}^*(n)DF_{p,m}^{*2}(n) = [m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))][m(F_{p,m}(n-1) + iF_{p,m}(n)) - \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))] = m^2[F_{p,m}^2(n-1) - F_{p,m}^2(n)] + i2m^2F_{p,m}(n-1)F_{p,m}(n)$, by equations (2) and (4).

$DF_{p,m}^*(n)DF_{p,m}^{*3}(n) = [m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))][m(F_{p,m}(n-1) - iF_{p,m}(n)) - \varepsilon(F_{p,m}(n-p-1) - iF_{p,m}(n-p))] = m^2[F_{p,m}^2(n-1) + F_{p,m}^2(n)] + i2m\varepsilon[F_{p,m}(n-p)F_{p,m}(n-1) - F_{p,m}(n)F_{p,m}(n-p-1)]$, by equations (2) and (5).

$$DF_{p,m}^*(n)DF_{p,m}^{*4}(n) = [m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))] [m(F_{p,m}(n-1) - iF_{p,m}(n))(1 - \varepsilon \frac{F_{p,m}(n-p-1) + iF_{p,m}(n-p)}{m(F_{p,m}(n-1) + iF_{p,m}(n))})] = m^2[F_{p,m}^2(n-1) + F_{p,m}^2(n)], \text{ by equation (2) and (6).}$$

$$DF_{p,m}^*(n)DF_{p,m}^{*5}(n) = [m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))] [m(F_{p,m}(n-p-1) + iF_{p,m}(n-p)) - \varepsilon(F_{p,m}(n-1) + iF_{p,m}(n))] = m^2[F_{p,m}(n-1)F_{p,m}(n-p-1) - F_{p,m}(n)F_{p,m}(n-p)] + im^2[F_{p,m}(n-1)F_{p,m}(n-p) + F_{p,m}(n)F_{p,m}(n-p-1)] + m\varepsilon[F_{p,m}^2(n-p-1) + F_{p,m}^2(n) - F_{p,m}^2(n-1) - F_{p,m}^2(n-p)] + i2m\varepsilon[F_{p,m}(n-p-1)F_{p,m}(n-p) - F_{p,m}(n-1)F_{p,m}(n)], \text{ by equations (2) and (7).}$$

$$DF_{p,m}^*(n) + DF_{p,m}^{*1}(n) = [m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))] + [m(F_{p,m}(n-1) - iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) - iF_{p,m}(n-p))] = 2[mF_{p,m}(n-1) + \varepsilon F_{p,m}(n-p-1)], \text{ by equations (2) and (3).}$$

$$DF_{p,m}^*(n) + DF_{p,m}^{*2}(n) = [m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))] + [m(F_{p,m}(n-1) + iF_{p,m}(n)) - \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))] = 2m[F_{p,m}(n-1) + iF_{p,m}(n)], \text{ by equations (2) and (4).}$$

$$DF_{p,m}^*(n) + DF_{p,m}^{*3}(n) = [m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))] + [m(F_{p,m}(n-1) - iF_{p,m}(n)) - \varepsilon(F_{p,m}(n-p-1) - iF_{p,m}(n-p))] = 2[mF_{p,m}(n-1) + i\varepsilon F_{p,m}(n-p)], \text{ by equations (2) and (5).}$$

$$(F_{p,m}(n-1) + iF_{p,m}(n))DF_{p,m}^{*4}(n) = (F_{p,m}(n-1) + iF_{p,m}(n)) [m(F_{p,m}(n-1) - iF_{p,m}(n))(1 - \varepsilon \frac{F_{p,m}(n-p-1) + iF_{p,m}(n-p)}{m(F_{p,m}(n-1) + iF_{p,m}(n))})] = F_{p,m}(n-1) - iF_{p,m}(n) [m(F_{p,m}(n-1) + iF_{p,m}(n)) - \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))] = [F_{p,m}(n-1) - iF_{p,m}(n)]DF_{p,m}^{*2}(n), \text{ by equations (4) and (6).}$$

$$\varepsilon DF_{p,m}^*(n) + mDF_{p,m}^{*5}(n) = m\varepsilon(F_{p,m}(n-1) + iF_{p,m}(n)) + m^2(F_{p,m}(n-p-1) + iF_{p,m}(n-p)) - m\varepsilon(F_{p,m}(n-1) + iF_{p,m}(n)) = m^2[F_{p,m}(n-p-1) + iF_{p,m}(n-p)], \text{ by equations (2) and (7).}$$

$$mDF_{p,m}^*(n) - \varepsilon DF_{p,m}^{*5}(n) = m[m(F_{p,m}(n-1) + iF_{p,m}(n)) + \varepsilon(F_{p,m}(n-p-1) + iF_{p,m}(n-p))] - \varepsilon[m(F_{p,m}(n-p-1) + iF_{p,m}(n-p)) - \varepsilon(F_{p,m}(n-1) + iF_{p,m}(n))] = m^2[F_{p,m}(n-1) + iF_{p,m}(n)], \text{ by equations (2) and (7).}$$

We know that $DF_{p,m}^*(n) = DF_{p,m}^*(n-1) + DF_{p,m}^*(n-p-1)$

We put $n = n + p + 1$, we get

$$DF_{p,m}^*(n+p+1) = DF_{p,m}^*(n+p) + DF_{p,m}^*(n) = DF_{p,m}^*(n) + DF_{p,m}^*(n+p).$$

Therefore, the norm of m -extension dual complex Fibonacci p -number, $NDF_{p,m}^*$ is defined in five different ways as follows:

$$\begin{aligned} NDF_{p,m}^{*1}(n) &= \|DF_{p,m}^*(n)DF_{p,m}^{*1}(n)\| \\ &= [m^2(F_{p,m}^2(n-1) + F_{p,m}^2(n)) + \\ &2m\varepsilon(F_{p,m}(n-1)F_{p,m}(n-p-1) + F_{p,m}(n)F_{p,m}(n-p))]^{\frac{1}{2}} \end{aligned}$$

$$\begin{aligned} NDF_{p,m}^{*2}(n) &= \|DF_{p,m}^*(n)DF_{p,m}^{*2}(n)\| \\ &= [(m^2(F_{p,m}^2(n-1) - F_{p,m}^2(n)))^2 + 4m^4F_{p,m}^2(n)F_{p,m}^2(n-1)]^{\frac{1}{2}} = \\ & m^2(F_{p,m}^2(n-1) + F_{p,m}^2(n)) \end{aligned}$$

$$NDF_{p,m}^{*3}(n) = \|DF_{p,m}^*(n)DF_{p,m}^{*3}(n)\| = m^2(F_{p,m}^2(n-1) + F_{p,m}^2(n))$$

$$\begin{aligned} NDF_{p,m}^{*4}(n) &= \|DF_{p,m}^*(n)DF_{p,m}^{*4}(n)\| = m(F_{p,m}^2(n-1) + F_{p,m}^2(n))^{\frac{1}{2}} \\ (NDF_{p,m}^{*5}(n))^2 &= (\|DF_{p,m}^*(n)DF_{p,m}^{*5}(n)\|)^2 = \\ m^4[F_{p,m}^2(n-1)F_{p,m}^2(n-p-1) + F_{p,m}^2(n)F_{p,m}^2(n-p) + F_{p,m}^2(n-1)F_{p,m}^2(n-p) \\ &+ F_{p,m}^2(n)F_{p,m}^2(n-p-1)] \\ &+ 2m^3\varepsilon[F_{p,m}(n-1)F_{p,m}^3(n-p-1) - F_{p,m}^3(n-1)F_{p,m}(n-p-1) + \\ &F_{p,m}(n)F_{p,m}^3(n-p) - F_{p,m}^3(n)F_{p,m}(n-p) + F_{p,m}(n-1)F_{p,m}(n-p-1)F_{p,m}^2(n-p) \\ &- F_{p,m}^2(n)F_{p,m}(n-1)F_{p,m}(n-p-1) + F_{p,m}(n)F_{p,m}(n-p)F_{p,m}^2(n-p-1) \\ &- F_{p,m}^2(n-1)F_{p,m}(n)F_{p,m}(n-p)]. \end{aligned}$$

□

Theorem 2.7. Let $n_1, n_2 \geq 0$ the Honsberger identity for m -extension dual complex Fibonacci p -numbers $DF_{p,m}^*(n_1), DF_{p,m}^*(n_2)$ is given by $DF_{p,m}^*(n_1)DF_{p,m}^*(n_2) + DF_{p,m}^*(n_1+1)DF_{p,m}^*(n_2+1) = m^2[F_{p,m}(n_1-1)F_{p,m}(n_2-1) - F_{p,m}(n_1+1)F_{p,m}(n_2+1)] + im^2[F_{p,m}(n_1-1)F_{p,m}(n_2) + F_{p,m}(n_1)F_{p,m}(n_2-1) + F_{p,m}(n_1+1)F_{p,m}(n_2) + F_{p,m}(n_1)F_{p,m}(n_2+1)] + \varepsilon m[F_{p,m}(n_1-1)F_{p,m}(n_2-p-1) + F_{p,m}(n_1-p-1)F_{p,m}(n_2-1) - F_{p,m}(n_1+1)F_{p,m}(n_2-p+1) - F_{p,m}(n_1-p+1)F_{p,m}(n_2+1)] + i\varepsilon m[F_{p,m}(n_1)F_{p,m}(n_2-p-1) + F_{p,m}(n_1-1)F_{p,m}(n_2-p) + F_{p,m}(n_1-p)F_{p,m}(n_2-1) + F_{p,m}(n_1-p-1)F_{p,m}(n_2) + F_{p,m}(n_1+1)F_{p,m}(n_2-p) + F_{p,m}(n_1)F_{p,m}(n_2-p+1) + F_{p,m}(n_1-p)F_{p,m}(n_2+1) + F_{p,m}(n_1-p+1)F_{p,m}(n_2)]$.

$$\begin{aligned} \text{Proof. } DF_{p,m}^*(n_1)DF_{p,m}^*(n_2) + DF_{p,m}^*(n_1+1)DF_{p,m}^*(n_2+1) &= [m(F_{p,m}(n_1-1) \\ &+ iF_{p,m}(n_1)) + \varepsilon F_{p,m}(n_1-p-1) + i\varepsilon F_{p,m}(n_1-p)][m(F_{p,m}(n_2-1) + \\ &iF_{p,m}(n_2)) + \varepsilon F_{p,m}(n_2-p-1) + i\varepsilon F_{p,m}(n_2-p)] \\ &+ [m(F_{p,m}(n_1) + iF_{p,m}(n_1+1)) + \varepsilon F_{p,m}(n_1-p) + i\varepsilon F_{p,m}(n_1-p+1)] \end{aligned}$$

$$\begin{aligned}
& [m(F_{p,m}(n_2) + iF_{p,m}(n_2 + 1)) + \varepsilon F_{p,m}(n_2 - p) + i\varepsilon F_{p,m}(n_2 - p + 1)] \\
& = m^2[F_{p,m}(n_1 - 1)F_{p,m}(n_2 - 1) - F_{p,m}(n_1 + 1)F_{p,m}(n_2 + 1)] + im^2[F_{p,m}(n_1 - \\
& 1)F_{p,m}(n_2) + F_{p,m}(n_1)F_{p,m}(n_2 - 1) + F_{p,m}(n_1 + 1)F_{p,m}(n_2) + F_{p,m}(n_1)F_{p,m}(n_2 \\
& + 1)] \\
& + \varepsilon m[F_{p,m}(n_1 - 1)F_{p,m}(n_2 - p - 1) + F_{p,m}(n_1 - p - 1)F_{p,m}(n_2 - 1) - \\
& F_{p,m}(n_1 + 1)F_{p,m}(n_2 - p + 1) - F_{p,m}(n_1 - p + 1)F_{p,m}(n_2 + 1)] \\
& + i\varepsilon m[F_{p,m}(n_1)F_{p,m}(n_2 - p - 1) + F_{p,m}(n_1 - 1)F_{p,m}(n_2 - p) + F_{p,m}(n_1 - \\
& p)F_{p,m}(n_2 - 1) + F_{p,m}(n_1 - p - 1)F_{p,m}(n_2) + F_{p,m}(n_1 + 1)F_{p,m}(n_2 - p) + \\
& F_{p,m}(n_1)F_{p,m}(n_2 - p + 1) + F_{p,m}(n_1 - p)F_{p,m}(n_2 + 1) + F_{p,m}(n_1 - p + \\
& 1)F_{p,m}(n_2)]. \quad \square
\end{aligned}$$

Theorem 2.8. Let $n_1, n_2 \geq 0$ the D'Ocagne's identity for m -extension dual complex Fibonacci p -numbers $DF_{p,m}^*(n_1), DF_{p,m}^*(n_2)$ is given by $DF_{p,m}^*(n_1)DF_{p,m}^*(n_2 + 1) - DF_{p,m}^*(n_1 + 1)DF_{p,m}^*(n_2) = m^2[F_{p,m}(n_1 - 1)F_{p,m}(n_2) - F_{p,m}(n_1)F_{p,m}(n_2 + 1) + F_{p,m}(n_1 + 1)F_{p,m}(n_2) - F_{p,m}(n_1)F_{p,m}(n_2 - 1)] + im^2[F_{p,m}(n_1 - 1)F_{p,m}(n_2 + 1) - F_{p,m}(n_1 + 1)F_{p,m}(n_2 - 1)] + \varepsilon m[F_{p,m}(n_1 - 1)F_{p,m}(n_2 - p) - F_{p,m}(n_1)F_{p,m}(n_2 - p + 1) + F_{p,m}(n_1 - p - 1)F_{p,m}(n_2) - F_{p,m}(n_1 - p)F_{p,m}(n_2 + 1) - F_{p,m}(n_1)F_{p,m}(n_2 - p - 1) + F_{p,m}(n_1 + 1)F_{p,m}(n_2 - p) - F_{p,m}(n_1 - p)F_{p,m}(n_2 - 1) + F_{p,m}(n_1 - p + 1)F_{p,m}(n_2)] + i\varepsilon m[F_{p,m}(n_1 - 1)F_{p,m}(n_2 - p + 1) + F_{p,m}(n_1 - p - 1)F_{p,m}(n_2 + 1) - F_{p,m}(n_1 + 1)F_{p,m}(n_2 - p - 1) - F_{p,m}(n_1 - p + 1)F_{p,m}(n_2 - 1)].$

Proof. $DF_{p,m}^*(n_1)DF_{p,m}^*(n_2 + 1) - DF_{p,m}^*(n_1 + 1)DF_{p,m}^*(n_2) = [m(F_{p,m}(n_1 - 1) + iF_{p,m}(n_1)) + \varepsilon F_{p,m}(n_1 - p - 1) + i\varepsilon F_{p,m}(n_1 - p)][m(F_{p,m}(n_2) + iF_{p,m}(n_2 + 1)) + \varepsilon F_{p,m}(n_2 - p) + i\varepsilon F_{p,m}(n_2 - p + 1)] - [m(F_{p,m}(n_1) + iF_{p,m}(n_1 + 1)) + \varepsilon F_{p,m}(n_1 - p) + i\varepsilon F_{p,m}(n_1 - p + 1)][m(F_{p,m}(n_2 - 1) + iF_{p,m}(n_2)) + \varepsilon F_{p,m}(n_2 - p - 1) + i\varepsilon F_{p,m}(n_2 - p)]$
 $= m^2[F_{p,m}(n_1 - 1)F_{p,m}(n_2) - F_{p,m}(n_1)F_{p,m}(n_2 + 1) + F_{p,m}(n_1 + 1)F_{p,m}(n_2) - F_{p,m}(n_1)F_{p,m}(n_2 - 1)] + im^2[F_{p,m}(n_1 - 1)F_{p,m}(n_2 + 1) - F_{p,m}(n_1 + 1)F_{p,m}(n_2 - 1)] + \varepsilon m[F_{p,m}(n_1 - 1)F_{p,m}(n_2 - p) - F_{p,m}(n_1)F_{p,m}(n_2 - p + 1) + F_{p,m}(n_1 - p - 1)F_{p,m}(n_2) - F_{p,m}(n_1 - p)F_{p,m}(n_2 + 1) - F_{p,m}(n_1)F_{p,m}(n_2 - p - 1) + F_{p,m}(n_1 + 1)F_{p,m}(n_2 - p) - F_{p,m}(n_1 - p)F_{p,m}(n_2 - 1) + F_{p,m}(n_1 - p + 1)F_{p,m}(n_2)] + i\varepsilon m[F_{p,m}(n_1 - 1)F_{p,m}(n_2 - p + 1) + F_{p,m}(n_1 - p - 1)F_{p,m}(n_2 + 1) - F_{p,m}(n_1 + 1)F_{p,m}(n_2 - p - 1) - F_{p,m}(n_1 - p + 1)F_{p,m}(n_2 - 1)]. \quad \square$

Theorem 2.9. Let $DF_{p,m}^*(n)$ be m -extension dual complex Fibonacci p -number. For $n \geq 1$, Cassini's identity for $DF_{p,m}^*(n)$ is given by $DF_{p,m}^*(n - 1)DF_{p,m}^*(n + 1) - D^2F_{p,m}^*(n) = m^2[F_{p,m}^2(n) - F_{p,m}^2(n - 1) + F_{p,m}(n - 2)F_{p,m}(n) - F_{p,m}(n + 1)F_{p,m}(n - 1)] + im^2[F_{p,m}(n + 1)F_{p,m}(n - 2) - F_{p,m}(n)F_{p,m}(n - 1)]$

$$\begin{aligned}
 & +\varepsilon m[F_{p,m}(n-2)F_{p,m}(n-p)-F_{p,m}(n-1)F_{p,m}(n-p+1)+F_{p,m}(n)F_{p,m}(n-p-2) \\
 & - F_{p,m}(n-p-1)F_{p,m}(n+1)-2F_{p,m}(n-1)F_{p,m}(n-p-1)+2F_{p,m}(n)F_{p,m}(n-p)] \\
 & +i\varepsilon m[F_{p,m}(n-2)F_{p,m}(n-p+1)-F_{p,m}(n-1)F_{p,m}(n-p)+F_{p,m}(n-p-2)F_{p,m}(n+1) \\
 & - F_{p,m}(n-p-1)F_{p,m}(n)].
 \end{aligned}$$

Proof. $DF_{p,m}^*(n-1)DF_{p,m}^*(n+1)-D^2F_{p,m}^*(n)$

$$\begin{aligned}
 & = [m(F_{p,m}(n-2)+iF_{p,m}(n-1))+\varepsilon(F_{p,m}(n-p-2)+iF_{p,m}(n-p-1))] \\
 & [m(F_{p,m}(n)+iF_{p,m}(n+1))+\varepsilon(F_{p,m}(n-p)+iF_{p,m}(n-p+1))] \\
 & - [m(F_{p,m}(n-1)+iF_{p,m}(n))+\varepsilon(F_{p,m}(n-p-1)+iF_{p,m}(n-p))]^2 \\
 & = m^2[F_{p,m}^2(n)-F_{p,m}^2(n-1)+F_{p,m}(n-2)F_{p,m}(n)-F_{p,m}(n+1)F_{p,m}(n-1)] \\
 & +im^2[F_{p,m}(n+1)F_{p,m}(n-2)-F_{p,m}(n)F_{p,m}(n-1)] \\
 & +\varepsilon m[F_{p,m}(n-2)F_{p,m}(n-p)-F_{p,m}(n-1)F_{p,m}(n-p+1)+F_{p,m}(n)F_{p,m}(n-p-2) \\
 & - F_{p,m}(n-p-1)F_{p,m}(n+1)-2F_{p,m}(n-1)F_{p,m}(n-p-1)+2F_{p,m}(n)F_{p,m}(n-p)] \\
 & +i\varepsilon m[F_{p,m}(n-2)F_{p,m}(n-p+1)-F_{p,m}(n-1)F_{p,m}(n-p)+F_{p,m}(n-p-2)F_{p,m}(n+1) \\
 & - F_{p,m}(n-p-1)F_{p,m}(n)]. \quad \square
 \end{aligned}$$

Theorem 2.10. *Let $DF_{p,m}^*(n+r)$ be the m -extension dual complex Fibonacci p -number. For $n \geq 1$, Catalan's identity for $DF_{p,m}^*(n+r)$ is given by $DF_{p,m}^*(n+r-1)DF_{p,m}^*(n+r+1)-D^2F_{p,m}^*(n+r)=m^2[F_{p,m}^2(n+r)-F_{p,m}^2(n+r-1)+F_{p,m}(n+r-2)F_{p,m}(n+r)-F_{p,m}(n+r+1)F_{p,m}(n+r-1)]+im^2[F_{p,m}(n+r+1)F_{p,m}(n+r-2)-F_{p,m}(n+r)F_{p,m}(n+r-1)]+\varepsilon m[F_{p,m}(n+r-2)F_{p,m}(n+r-p)-F_{p,m}(n+r-1)F_{p,m}(n+r-p+1)+F_{p,m}(n+r)F_{p,m}(n+r-p-2)-F_{p,m}(n+r-p-1)F_{p,m}(n+r+1)-2F_{p,m}(n+r-1)F_{p,m}(n+r-p-1)+2F_{p,m}(n+r)F_{p,m}(n+r-p)]+i\varepsilon m[F_{p,m}(n+r-2)F_{p,m}(n+r-p+1)-F_{p,m}(n+r-1)F_{p,m}(n+r-p)+F_{p,m}(n+r-p-2)F_{p,m}(n+r+1)-F_{p,m}(n+r-p-1)F_{p,m}(n+r)]$.*

Proof. $DF_{p,m}^*(n+r-1)DF_{p,m}^*(n+r+1)-D^2F_{p,m}^*(n+r)=$

$$\begin{aligned}
 & [m(F_{p,m}(n+r-2)+iF_{p,m}(n+r-1))+\varepsilon(F_{p,m}(n+r-p-2)+iF_{p,m}(n+r-p-1))] \\
 & [m(F_{p,m}(n+r)+iF_{p,m}(n+r+1))+\varepsilon(F_{p,m}(n+r-p)+iF_{p,m}(n+r-p+1))] \\
 & - [m(F_{p,m}(n+r-1)+iF_{p,m}(n+r))+\varepsilon(F_{p,m}(n+r-p-1)+iF_{p,m}(n+r-p))]^2 \\
 & = m^2[F_{p,m}^2(n+r)-F_{p,m}^2(n+r-1)+F_{p,m}(n+r-2)F_{p,m}(n+r)-F_{p,m}(n+r+1)F_{p,m}(n+r-1)] \\
 & +im^2[F_{p,m}(n+r+1)F_{p,m}(n+r-2)-F_{p,m}(n+r)F_{p,m}(n+r-1)] \\
 & +\varepsilon m[F_{p,m}(n+r-2)F_{p,m}(n+r-p)-F_{p,m}(n+r-1)F_{p,m}(n+r-p+1)+F_{p,m}(n+r)F_{p,m}(n+r-p-2) \\
 & - F_{p,m}(n+r-p-1)F_{p,m}(n+r+1)-2F_{p,m}(n+r-1)F_{p,m}(n+r-p-1)+2F_{p,m}(n+r)F_{p,m}(n+r-p)] \\
 & +i\varepsilon m[F_{p,m}(n+r-2)F_{p,m}(n+r-p+1)-F_{p,m}(n+r-1)F_{p,m}(n+r-p)+F_{p,m}(n+r-p-2)F_{p,m}(n+r+1) \\
 & - F_{p,m}(n+r-p-1)F_{p,m}(n+r)]. \quad \square
 \end{aligned}$$

3. CONCLUSION

In this paper, we introduced m -extension dual complex Fibonacci p -numbers. We established the properties of m -extension dual complex Fibonacci p -numbers. We also established the Honsberger identity, D'Ocagne's identity, Cassini's identity and Catalan's identity for m -extension dual complex Fibonacci p -numbers. I hope that these results will be useful in applied mathematics, quantum mechanics, quantum physics, Lie groups, number theory, Kinematics and differential equations like dual complex k -Fibonacci numbers [1], dual complex Fibonacci p -numbers [14], dual-complex numbers and their holomorphic functions [11] and dual-complex Jacobsthal quaternions [2] etc.

Acknowledgments

The author would like to thank the editor and anonymous reviewers for their valuable comments and suggestions to improve the quality of the paper.

REFERENCES

1. F. T. Aydin, *Dual Complex k -Fibonacci numbers*, Chaos Solitons Fractals, **115** (2018), 1-6.
2. F. T. Aydin, *Dual-complex Jacobsthal Quaternions*, Mathematical Sciences and Applications E-Notes, **8(2)** (2020), 145-156.
3. M. Basu and B. Prasad, *The generalized relations among the code elements for Fibonacci coding theory*, Chaos Solitons Fractals, **41** (2009), 2517-2525.
4. M. Basu and B. Prasad, *Long range variations on the Fibonacci universal code*, J. Number Theory, **130** (2010), 1925-1931.
5. W. K. Clifford, *Preliminary sketch biquaternions*, Proc London Math Soc, (4) **64** (1873), 381-395.
6. M. S. EL Naschie, *Topics in the mathematical physics of E-infinity theory*, Chaos Solitons Fractals, **30** (2006), 656-663.
7. M. S. EL Naschie, *The theory of cantor space time and high energy particle physics*, Chaos Solitons Fractals, **41** (2009), 2635-2646.
8. M. A. Gungor and A.Z. Azak, *Investigation of dual complex Fibonacci, dual complex Lucas numbers and their properties*, Adv Appl Clifford Alg, (4) **27** (2017), 3083-3096.
9. E. Kocer EG et al, *On the m extension of the Fibonacci and Lucas p -numbers*, Chaos Solitons Fractals, (4) **40** (2009), 1890-1906.
10. A. Kotelnikov, *Screw calculus and some of its applications to geometry and mechanics*, Annals of Imperial University of Kazan 1985.
11. F. Messelmi, *Dual complex numbers and their holomorphic functions*, 2015. Working paper or preprint: URL [https:// hal.archives-ouvertes.fr/hal-01114178](https://hal.archives-ouvertes.fr/hal-01114178).

12. B. Prasad, *The generalized relations among the code elements for a new complex Fibonacci matrix*, Discrete Math. Algorithms Appl. (2) **11** (2019), 1950026 (16 pages).
13. B. Prasad, *A new Gaussian Fibonacci matrices and its applications*, J. Algebra Relat. Topics, (1) **7** (2019), 65-72.
14. B. Prasad, *Dual Complex Fibonacci numbers p -numbers*, Chaos Solitons Fractals, **145** (2021), 109922.
15. A. P. Stakhov, *Fibonacci matrices, a generalization of the cassini formula and a new coding theory*, Chaos Solitons Fractals, **30** (2006), 56-66.
16. A. P. Stakhov, *The golden matrices and a new kind of cryptography*, Chaos Solitons Fractals, **32** (2007), 1138-1146.
17. E. Study, *Geometrie der dynamen*, Leipzig 1903.

Bandhu Prasad

Department of Mathematics, Kandi Raj College, Kandi, India

Email: bandhu_iit@rediffmail.com