

Join maximal element graph of lattice module

Lakpa Sherpa[†], Vilas Kharat[†], Manish Agalave[‡], Narayan Phadatare^{§*}

[†]Department of Mathematics, Savitribai Phule Pune University, Pune, India

[‡]Department of Mathematics, Fergusson College(Autonomus), Pune, India

[§] Bharati Vidyapeeth Deemed to be University College of Engineering, Pune, India

Emails: csherpaap@gmail.com, laddoo1@yahoo.com, manishagalave@gmail.com,
a9999phadatare@gmail.com

Abstract. Let \mathcal{L} be a C -lattice and M be a lattice module over \mathcal{L} . The join maximal element graph $\mathbb{G}(M)$ is a simple, undirected graph with all proper non-zero elements of M as vertices, and two distinct vertices, N and K , are adjacent if and only if $N \vee K \in \text{Max}(M)$, where $\text{Max}(M)$ is the collection of all maximal elements of M . In this paper, some properties of the graph $\mathbb{G}(M)$ like diameter, girth and clique number are investigated. Also, the interplay between the algebraic properties of M and the properties of those graphs is studied.

Keywords: Maximal element, Jacobson radical $J_{rad}(M)$, Join maximal element graph $\mathbb{G}(M)$.

AMS Subject Classification 2010: 06B23, 05C99, 06E10, 06F99.

1 Introduction

The zero-divisor graphs of a commutative ring and its various generalizations has been studied in ([6–11], [13, 14] and [22]). In 1988, I. Beck defined the concept of zero-divisor graphs with the help of commutative rings (see [12]). This study is further extended by Akbari et al. [3] and introduced the intersection graph of a R -module M . In [5], A. H. Alwan generalized the concept of maximal ideal graph of a ring (see [4]) and defined the new graph called as maximal submodule graph of a module.

Various graph structures associated with module over ring structure can be identified in ([1], [23]). In this paper, we used the concept introduced by A. H. Alwan to define the join maximal element graph of \mathcal{L} -module M , and throughout this paper \mathcal{L} is a multiplicative lattice and M is a lattice module over \mathcal{L} .

A *multiplicative lattice* is denoted as $(\mathcal{L}, 0_{\mathcal{L}}, 1_{\mathcal{L}}, *)$, where \mathcal{L} is a complete lattice with least element $0_{\mathcal{L}}$, greatest element $1_{\mathcal{L}}$ and $*$ is a binary operation defined on \mathcal{L} that satisfies the

*Corresponding author

Received: 27 August 2024/ Revised: 23 April 2025/ Accepted: 10 February 2026

DOI: [10.22124/JART.2026.28291.1704](https://doi.org/10.22124/JART.2026.28291.1704)

following properties for all $a, b, c \in \mathcal{L}$:

1. $a * b \leq a \wedge b$.
2. $a * b = b * a$.
3. $(a * b) * c = a * (b * c)$.
4. $a * (\bigvee_{\alpha \in I} b_\alpha) = \bigvee_{\alpha \in I} (a * b_\alpha)$, where $b_\alpha \in \mathcal{L}$ and I is an indexing set.
5. $a * 1_{\mathcal{L}} = a$.

Henceforth, we write $a * b = ab$ for convenience only. By a C -lattice we mean a multiplicative lattice $(\mathcal{L}, 0_{\mathcal{L}}, 1_{\mathcal{L}}, *)$ in which the greatest element $1_{\mathcal{L}}$ is compact as well as multiplicative identity and $(\mathcal{L}, 0_{\mathcal{L}}, 1_{\mathcal{L}}, *)$ is generated under join by a multiplicatively closed set C of compact elements. For more reading about multiplicative lattices, one may refer to ([17], [18]- [19]).

Let \mathcal{L} be a multiplicative lattice. A *lattice module* or an \mathcal{L} -*module* is complete lattice M with least elements 0_M and the greatest element 1_M over \mathcal{L} if for $u \in \mathcal{L}$ and $N \in M$ a multiplication denoted by $u.N \in M$ satisfies the following properties:

1. $(uv).N = u.(v.N)$;
2. $(\bigvee_{\alpha} u_{\alpha}).(\bigvee_{\beta} N_{\beta}) = (\bigvee_{\alpha\beta} u_{\alpha}.N_{\beta})$;
3. $1_{\mathcal{L}}.N = N$;
4. $0_{\mathcal{L}}.N = 0_M$; for all $u, v, u_{\alpha} \in \mathcal{L}$, and for all $N, N_{\beta} \in M$.

An element $N \in M$ is called *meet-principal*, if for each $a \in \mathcal{L}$ and $A \in M$, we have $A \wedge aN = (a \wedge (A : N))N$. An element $N \in M$ is called *join-principal*, if $((aN \vee A) : N) = (a \vee (A : N))$ for each $a \in \mathcal{L}$ and $A \in M$. If N is both meet-principal and join-principal, then N is called *principal* element of M . Note that, if each member of M is the join of principal elements then M is called *principally generated lattice* or *PG-lattice*.

Throughout this paper, $u.N$ will be written as uN . For $N \in M$ and $b \in \mathcal{L}$, denote $(N : b) = \bigvee\{K \in M \mid bK \leq N\}$ and for $A, B \in M$, $(A : B) = \bigvee\{x \in \mathcal{L} \mid xB \leq A\}$. Here the operation $:$ is called *residual division*.

A proper element $N \in M$ is a *maximal element* if for every element $B \in M$ such that $N \leq B$, implies either $N = B$ or $B = 1_M$. Denote $Max(M)$ for the collection of all maximal elements of a lattice module M over a C -lattice \mathcal{L} . The meet of all members of $Max(M)$ is called the *Jacobson radical* of M and it is denoted by $J_{rad}(M)$.

For various concepts and related aspects of the lattice module, we refer ([2], [16], [18] and [21]).

Let $G = (V, E)$ be a graph with the set of vertex $V = V(G)$ and the set of edges $E = E(G)$. If $E = \emptyset$, then G is called an *empty graph*. Let $b \in V$, the *degree* of b in a graph G refers to the number of edges incident to that vertex and it is denoted by $deg(b)$. For $a, b \in V$, the length of the shortest path between a and b is denoted by $d(a, b)$. If there is no path between a and b , then $d(a, b) = \infty$. The *diameter* of G is $dim(G) = \sup\{d(a, b) \mid a, b \in V\}$. A graph's greatest complete subgraph is called a *clique*. The number of vertices in largest clique is called the clique number of G , which is represented as $\omega(G)$. The *girth* of graph G , denoted by $gr(G)$, is the

length of the shortest cycle in G . For $a \in V$, the eccentricity $e(a)$ of a is the greatest distance between a and any other vertex b in G , i.e., $e(a) = \text{Max}\{d(a,b) \mid b \in V\}$. Let G be a connected graph. The *radius* of G , denoted by $\zeta(G)$, is essentially the minimum eccentricity among all the vertices in G , i.e., $\zeta(G) = \text{Min}\{e(u) \mid u \in V\}$. For $a \in V$, the collection $\{b_i\}$ of all adjacent vertices to a is called *neighborhood* of a , and it is denoted by $\Delta(a)$.


For more information on graph theory, the reader may refer ([15], [20], [24]).

In this paper, we prove that the graph $\mathbb{G}(M)$ is not a complete graph if the lattice module M has more than one maximal elements. Also, we show that the number of maximal elements of M can not be greater than 5 if the graph $\mathbb{G}(M)$ is planar. It is proved that, the graph $\mathbb{G}(M)$ is connected with $\text{dim}(\mathbb{G}(M)) \leq 4$, under certain conditions.

2 Connectivity, diameter and girth of $\mathbb{G}(M)$


Definition 1. Let M be a \mathcal{L} -module. The join maximal element graph of M , which is denoted as $\mathbb{G}(M)$ is simple, undirected graph with all proper non-zero elements of M as vertices and two distinct vertices N and K are adjacent if and only if $N \vee K \in \text{Max}(M)$.

Example 1. The lattice depicted in Fig.(4.1) is a C -lattice \mathcal{L} and the lattice depicted in Fig.(4.2) is a lattice module M over \mathcal{L} . Fig.(4.3) represents the join maximal element graph $\mathbb{G}(M)$ of M as shown in Fig.(4.2) with vertex set $V(\mathbb{G}(M)) = \{A, B, C, Q\}$.




.	0	a	b	c	d	$1_{\mathcal{L}}$
0	0	0	0	0	0	0
a	0	a	0	a	0	a
b	0	0	0	0	b	b
c	0	a	0	a	b	c
d	0	0	b	b	d	d
$1_{\mathcal{L}}$	0	a	b	c	d	$1_{\mathcal{L}}$

Fig.(4.1) Multiplicative Lattice \mathcal{L}



.	0_M	A	B	C	Q	1_M
0	0_M	0_M	0_M	0_M	0_M	0_M
a	0_M	A	0_M	A	0_M	A
b	0_M	0_M	0_M	0_M	B	B
c	0_M	A	0_M	A	B	C
d	0_M	0_M	B	B	Q	Q
$1_{\mathcal{L}}$	0_M	A	B	C	Q	1_M

Fig.(4.2) Lattice Module M over itself

Fig.(4.3) Graph $\mathbb{G}(M)$ of M

Note that, each $N \notin \text{Max}(M)$ is adjacent to at least one $K \in \text{Max}(M)$. Also, for each $K_i \in \text{Max}(M)$, $J_{\text{rad}}(M)$ is adjacent to K_i . We know that for two distinct elements $N, K \in M$, if $N \vee K = 1_M$, then N and K are said to be *co-maximal*. It is clear that the co-maximal elements of M are not adjacent in $\mathbb{G}(M)$ and so have the following.

Throughout the paper, we fundamentally need the following Lemma 1.

Lemma 1 ([21]). *Let M be a lattice module over a C -lattice L with 1_M is compact. Then for every proper element $X \in M$, there exists a maximal element N of M such that $X \leq N$.*

Theorem 1. *Let M be a \mathcal{L} -module with 1_M compact and $1 < |\text{Max}(M)| < \infty$. Then there is no vertex in $V(\mathbb{G}(M))$ which is adjacent to the remaining vertices. Furthermore, the graph $\mathbb{G}(M)$ is not a complete graph.*

Theorem 2. *Let M be a \mathcal{L} -module with 1_M compact and $N, P, K \in V(\mathbb{G}(M))$ with $K \in \text{Max}(M)$. Then the following statements hold:*

1. $K \in \Delta(N) \cap \Delta(P)$ if and only if $K \in \Delta(N \vee P)$, where $N \vee P \neq K, 1_M$.
2. If $N \leq J_{\text{rad}}(M)$, then $K \in \Delta(N)$.
3. If $N \leq P$ and $P \notin \text{Max}(M)$, then $N \notin \Delta(P)$.

Proof. 1) Suppose that $K \in \Delta(N) \cap \Delta(P)$. Then by definition, K is adjacent to N and P . Since $K \in \text{Max}(M)$, we have $N \leq K$ and $P \leq K$, and hence $N \vee P \leq K$. This implies that $K \in \Delta(N \vee P)$. The converse part is clear.

2) Let $N \leq J_{\text{rad}}(M)$. Then $N \leq J_{\text{rad}}(M) \leq K$. This implies that $N \leq K$ and hence $K \in \Delta(N)$.

3) If $N \leq P$, then $N \vee P = P$. But $P \notin \text{Max}(M)$, therefore N is not adjacent to P . \square

Theorem 3. *Let M be a \mathcal{L} -module with 1_M compact and $\mathbb{G}(M)$ be the join maximal element graph of M . If $P, Q \notin \text{Max}(M)$ are adjacent vertices, then there exists only one $K \in \text{Max}(M)$ such that $K \in \Delta(P) \cap \Delta(Q)$.*

Proof. Suppose that $K_1, K_2 \in \text{Max}(M)$ such that $K_1, K_2 \in \Delta(P) \cap \Delta(Q)$. By definition, P, Q are adjacent to both K_1, K_2 in $\mathbb{G}(M)$. Since P, Q are adjacent vertices, we have $P \vee Q \in \text{Max}(M)$, therefore $K_1 = P \vee Q = K_2$. \square

Theorem 4. *Let M be a \mathcal{L} -module with 1_M compact and $J_{\text{rad}}(M) \notin \text{Max}(M) \cup \{0_M\}$. Then $\mathbb{G}(M) \cong K_{m,n}$, where $m, n \in \mathbb{Z}^+$ if and only if every member of $M \setminus \text{Max}(M)$ (i.e., non maximal element) is less than $J_{\text{rad}}(M)$.*

Proof. Suppose that every member of $M \setminus \text{Max}(M)$ is less than $J_{\text{rad}}(M)$. Let $V_1 = \text{Max}(M)$ and $V_2 = \{P \in V(\mathbb{G}(M)) \mid P \leq J_{\text{rad}}(M)\}$. By Remark 2, any two vertices $P, Q \in V_1$ are not adjacent in $\mathbb{G}(M)$. Since $J_{\text{rad}}(M) \notin \text{Max}(M)$, for any $P, K \in V_2$ we have $P \vee K \notin \text{Max}(M)$. This implies that any two vertices of V_2 are not adjacent in $\mathbb{G}(M)$. Also, by Theorem 2(2), it is observed that each $N \in V_1$ is adjacent to each $K \in V_2$ and hence $\mathbb{G}(M) \cong K_{m,n}$, where $m = |V_1|$ and $n = |V_2|$. Conversely, if $\mathbb{G}(M)$ is a complete bipartite graph with two part Y_1 and Y_2 . It is very clear that $Y_1 = \text{Max}(M)$ and $Y_2 = \{P \in V(\mathbb{G}(M)) \mid P \leq J_{\text{rad}}(M)\}$. \square

Corollary 1. *Let M be a \mathcal{L} -module with 1_M compact and $J_{\text{rad}}(M) \notin \text{Max}(M) \cup \{0_M\}$. If $\mathbb{G}(M)$ is not complete bipartite graph and $|V(\mathbb{G}(M)) \setminus (\text{Max}(M) \cup \{P \in V(\mathbb{G}(M)) \mid P \leq J_{\text{rad}}(M)\})| = 1$, then $\mathbb{G}(M)$ is tripartite.*

Proof. Since $\mathbb{G}(M)$ is not complete bipartite graph, by Theorem 4 we have $P \not\leq J_{\text{rad}}(M)$ for some $P \in M \setminus \text{Max}(M)$. Now, let $V_1 = \text{Max}(M)$, $V_2 = \{P \in V(\mathbb{G}(M)) \mid P \leq J_{\text{rad}}(M)\}$ and $V_3 = V(\mathbb{G}(M)) - V_1 \cup V_2$. It is simple to demonstrate that each pair of vertices in V_i is non adjacent for $i = 1, 2, 3$. Hence, V_1, V_2 and V_3 are nothing but 3 partition of $\mathbb{G}(M)$. \square

A *star* graph is a particular kind of graph wherein nodes are connected to other nodes, referred to as leaves, by edges, starting at the center node. Although the leaf nodes are not connected to one another, every leaf node is connected to the central node.

Proposition 1. *Let M be a \mathcal{L} -module with 1_M compact. If $M = C_n$, i.e., M is a chain with n elements, then $\mathbb{G}(M)$ is a star graph.*

Proof. Proof is straightforward. \square

Theorem 5. *Let M be a \mathcal{L} -module with 1_M compact and $J_{\text{rad}}(M) \neq 0_M$. Then $gr(\mathbb{G}(M))$, is either 3 or 4 or ∞ .*

Proof. If there is no any cycle in $\mathbb{G}(M)$ then $gr(\mathbb{G}(M)) = \infty$. Suppose that vertex $P, K \in V(\mathbb{G}(M)) \setminus \text{Max}(M)$ are adjacent in $\mathbb{G}(M)$. Then by definition, $P, K \neq P \vee K \in \text{Max}(M)$. Therefore we have a cycle $P - K - P \vee K - P$ of length 3 in $\mathbb{G}(M)$. Now suppose that for every adjacent vertices X, Y either $X \in \text{Max}(M)$ or $Y \in \text{Max}(M)$ and let $N_1 - N_2 - \dots - N_{n-1} - N_n - N_1$ be a cycle of length n . By Remark 2, members of $\text{Max}(M)$ are not adjacent in $\mathbb{G}(M)$, therefore the vertices in cycle $N_1 - N_2 - \dots - N_{n-1} - N_n - N_1$ are alternatively member of $\text{Max}(M)$ and non-member of $\text{Max}(M)$ and hence $J_{\text{rad}}(M) \notin \text{Max}(M)$. Let $N_1 \in \text{Max}(M)$. Then $J_{\text{rad}}(M)$ is adjacent to maximal elements in cycle $N_1 - N_2 - \dots - N_{n-1} - N_n - N_1$, say N_1, N_3, N_5, \dots . If $J_{\text{rad}}(M) = N_2$, then we have a cycle $J_{\text{rad}}(M) = N_2 - N_3 - N_4 - N_5 - N_2$. If $J_{\text{rad}}(M) \neq N_2$, then $J_{\text{rad}}(M) - N_1 - N_2 - N_3 - J_{\text{rad}}(M)$ is a cycle in the graph $\mathbb{G}(M)$. From both the cases, it is observed that, $gr(\mathbb{G}(M))$ is either 3 or 4. \square

A graph G in which every pair of vertices is connected by a path is said to be a *connected* graph.

Theorem 6. *Let M be a \mathcal{L} -module with 1_M compact and $\mathbb{G}(M)$ be the join maximal element graph of M . If whenever $N \wedge K \neq 0_M$ for any two distinct element N, K of M , then $\mathbb{G}(M)$ is connected graph with $\dim(\mathbb{G}(M)) \leq 4$.*

Proof. Suppose that $N, K \in V(\mathbb{G}(M))$ and $N \neq K$. If $N - K$ then we are through. Suppose that N and K are not adjacent. This implies that either $N \vee K = 1_M$ or $N \vee K < P$ for some $P \in \text{Max}(M)$. For $N \vee K < P$, we have a path $N - P - K$ of length equal to 2 in $\mathbb{G}(M)$. Suppose that $N \vee K = 1_M$. This implies following cases:

(i) At least one of N and K is a member of $\text{Max}(M)$ and they are not comparable.

(ii) Both N and K are not member of $\text{Max}(M)$ and they are not comparable.

Suppose that case (i) holds and $N \in \text{Max}(M)$. If $K \in \text{Max}(M)$, then we have path $N - N \wedge K - K$ between N and K in $\mathbb{G}(M)$. Let $K \notin \text{Max}(M)$. Then we have $P \in \text{Max}(M)$ such that K and P are adjacent in the graph $\mathbb{G}(M)$. If $P = N$, then we have a path $K - N$ of length equal to 1 in $\mathbb{G}(M)$. If $P \neq N$, then $N - N \wedge P - P - K$ is a path in $\mathbb{G}(M)$. Thus in case (i), it is observed that N and K are connected and $\dim(\mathbb{G}(M)) \leq 3$.

If case (ii) holds. Since $N \wedge K \neq 0$, by Lemma 1 we have $X \in \text{Max}(M)$ such that $N \wedge K \leq X$. Thus $N - N' - N \wedge K - K' - K$ is a path of length equal to 4 in $\mathbb{G}(M)$. Consequently, from each case, we have any $N, K \in V(\mathbb{G}(M))$ that are connected and $\dim(\mathbb{G}(M)) \leq 4$. \square

A *cut vertex* is a vertex in a graph G such that if we remove it (along with all incident edges), the graph becomes disconnected or has at least two connected components. In other words, removing a cut vertex from the graph G increases the number of connected components.

Theorem 7. *Let M be a \mathcal{L} -module with 1_M compact and $N \wedge K \neq 0_M$ for any two distinct element N, K of M . If Y is a cut vertex of the graph $\mathbb{G}(M)$, then $Y = P \wedge Q$ for $P, Q \in \text{Max}(M)$.*

Proof. If $Y \in \text{Max}(M)$, then the proof is clear. Suppose that $Y \notin \text{Max}(M)$. Since Y is a cut vertex, we have different components in $\mathbb{G}(M) - \{Y\}$. Let X_1 and X_2 be two vertices in different components of $\mathbb{G}(M) - \{Y\}$. Consider the following cases:

Case-1) If $X_1, X_2 \in \text{Max}(M)$, then it is clearly observed that $X_1 \wedge X_2 \in \Delta(X_1) \cap \Delta(X_2)$ and hence $Y = X_1 \wedge X_2$.

Case-2) If $X_1 \in \text{Max}(M)$ and $X_2 \notin \text{Max}(M)$. Then we have $S \in \text{Max}(M)$ such that $X_2 \in \Delta(S)$. Since $X_1 \wedge S$ is adjacent to X_1 and S , therefore $Y = X_1 \wedge S$.

Case-3) If $X_1, X_2 \notin \text{Max}(M)$. Then there is $C, D \in \text{Max}(M)$ such that C and D are adjacent to X_1 and X_2 respectively. As Y is cut vertex, we should have $C \neq D$. Thus as like Case-2), we have $Y = C \wedge D$. \square

An *induced subgraph* of a graph G is a subgraph that is formed by selecting a subset of vertices from the original graph along with all the edges that are present between those vertices in the original graph. In other words, an induced subgraph retains all edges between the selected vertices and includes no additional edges.

Proposition 2. *Let M be a \mathcal{L} -module with 1_M compact and for unique $Z \in \text{Max}(M)$, $\Pi(M)$ be the induced subgraph by $\{K \in V(\mathbb{G}(M)) | K \leq Z\}$. Then the clique of $\mathbb{G}(M)$ is contained in $\Pi(M)$.*

Proof. Suppose that Γ is the clique of $\mathbb{G}(M)$. Since any two distinct maximal elements $N, P \in \text{Max}(M)$ are not adjacent in $\mathbb{G}(M)$, we have only one maximal element in Γ . Thus by definition of Γ and Proposition 3, we have unique maximal element $Z \in \text{Max}(M)$ such that Γ is a subgraph induced by $\{K \in V(\mathbb{G}(M)) | K \leq Z\}$. \square

Theorem 8. *Let M be a \mathcal{L} -module with 1_M compact and $J_{rad}(M) \neq \emptyset$. If $|Max(M)| \geq 2$ then $\zeta(\mathbb{G}(M)) = 2$.*

Proof. Since $J_{rad}(M)$ is adjacent to each $K_i \in Max(M)$, we have $d(J_{rad}(M), T) = 1$ for any $T \in Max(M)$. Also, since each $N \notin Max(M)$ is adjacent to at least one $K \in Max(M)$, we have $d(J_{rad}(M), K) \leq 2$ for any $K \notin Max(M)$. Let $X, Y \in Max(M)$ with $X \neq Y$. If $X \wedge Y$ is adjacent to $J_{rad}(M)$, then we have $P \in Max(M)$ such that $J_{rad}(M) \vee (X \wedge Y) = P$. But $J_{rad}(M) \vee (X \wedge Y) \leq X, Y$, therefore $P = X = Y$, which is contradiction to $X \neq Y$. This implies that $X \wedge Y$ is not adjacent to $J_{rad}(M)$. So the eccentricity e of $J_{rad}(M)$ is 2. Suppose that $F \in V(\mathbb{G}(M))$ with $e(F) = 1$. Then F is adjacent to each $W \in Max(M)$. It is clear that $F \notin Max(M)$. Since for any $X, Y \in Max(M)$ with $X \neq Y$, $X \wedge Y$ is not adjacent to $J_{rad}(M)$, we have neither $F = X \wedge Y$ nor $F = J_{rad}(M)$ and hence $F < J_{rad}(M)$. Thus F is adjacent to each $W \in Max(M)$ and $F < J_{rad}(M)$ which contradicts that $e(F) = 1$, therefore $J_{rad}(M)$ has the minimum eccentricity among all the vertices of $\mathbb{G}(M)$. Consequently, $\zeta(\mathbb{G}(M)) = 2$. \square

We will now investigate the planar property of the join maximal element graph $\mathbb{G}(M)$.

Proposition 3. *Let M be a \mathcal{L} -module with 1_M compact and $\mathbb{G}(M)$ be the join maximal element graph of M . If for each $Z \in Max(M)$, the induced subgraph by $\{K \in V(\mathbb{G}(M)) | K \leq Z\}$ is planar, then $\omega(\mathbb{G}(M)) \in \{2, 3, 4\}$.*

Proof. Proof follows from Proposition 2 and Koratowski's theorem [15]. \square

In the following theorem we prove that the planarity of the join maximal element graph $\mathbb{G}(M)$ put the control on $|Max(M)|$.

Theorem 9. *Let M be a \mathcal{L} -module with 1_M compact and $J_{rad}(M) \neq \emptyset$. If the join maximal element graph $\mathbb{G}(M)$ is planar, then $|Max(M)| \leq 4$.*

Proof. Suppose that $\mathbb{G}(M)$ is planar graph and the \mathcal{L} -module M has at least five maximal elements say P, Q, R, S and T . Then it is clear that any one element from the set $\{P \wedge Q \wedge R, P \wedge Q \wedge R \wedge S, P \wedge Q \wedge R \wedge S \wedge T\}$ is non-zero vertex which is adjacent to each of P, Q and R in $\mathbb{G}(M)$. This implies that the graph $\mathbb{G}(M)$ contains $K_{3,3}$, i.e., complete bipartite graph, which contradicts the fact that $\mathbb{G}(M)$ is planar graph. Consequently, $|Max(M)| \leq 4$. \square

Acknowledgments

The authors express their sincere gratitude to the reviewers and editors for investing their time and effort in providing insightful suggestions and comments, which significantly enhanced the quality of work.

References

- [1] A. Abbasi and Sh. Habibi, *The total graph of a module over a commutative ring with respect to proper submodules*, J. Algebra Appl., (3) **11** (2012), 1250048.

- [2] E. A. AL-Khouja, *Maximal elements and prime elements in lattice modules*, Damascus Univ. Basic Sci., **19** (2003), 9-20.
- [3] S. Akbari, H. A. Tavallaee, and S. Khalashi Ghezelahmad, *Intersection graph of submodules of a module*, J. Algebra Appl., (1) **11** (2012), 1250019.
- [4] F. H. Abdulqadr, *Maximal ideal graph of commutative rings*, Iraqi J. Sci., (8) **61** (2020), 2070-2076.
- [5] A. H. Alwan, *Maximal submodule graph of a module*, J. Discrete Math. Sci. Cryptogr., (7) **24** (2021), 1941-1949.
- [6] S. Akbari and A. Mohammadian, *On the zero-divisor graph of a commutative ring*, J. Algebra, (2) **274** (2004), 847-855.
- [7] D. F. Anderson, R. Levy and J. Shapiro, *Zero-divisor graphs, von Neumann regular rings and Boolean algebras*, J. Pure Appl. Algebra, **180** (2003), 221-241.
- [8] F. F. Anderson and M. Naseer, *Beck's coloring of a commutative ring*, J. Algebra, **159** (1993), 500-514.
- [9] D. F. Anderson and P. S. Livingston, *The zero-divisor graph of a commutative ring*, J. Algebra, (2) **217** (1999), 434-447.
- [10] H. Ansari-Toroghy and Sh. Habibi, *The zariski topology-graph of modules over commutative rings*, Commun. Algebra, **42** (2014), 3283-3296.
- [11] H. Ansari-Toroghy, F. Farshadifar and F. Mahboobi-Abkenar, *The small intersection graph relative to multiplication modules*, J. Algebra Relat. Topics, (1) **4** (2016), 21-32.
- [12] I. Beck, *Coloring of commutative rings*, J. Algebra, **116** (1988), 208-226.
- [13] M. Behboodi and Z. Rakeei, *The annihilating-ideal graph of commutative rings-I*, J. Algebra Appl., (4) **10** (2011), 727-739.
- [14] M. Behboodi and Z. Rakeei, *The annihilating-ideal graph of commutative rings-II*, J. Algebra Appl., (4) **10** (2011), 741-753.
- [15] J. Bondy and U. Murty, *Graph Theory, Graduate Texts in Mathematics 244*, Springer, New York, 2008.
- [16] F. Callialp, G. Ulucak and U. Tekir, *On the Zariski topology over an L -module M* , Turk. J. Math., **41** (2017), 326-336.
- [17] F. Callialp, C. Jayaram and U. Tekir, *Weakly prime elements in multiplicative lattices*, Commun. Algebra, **40** (2012), 2825-2840.
- [18] J. A. Johnson, *a -Adic completions of Noetherian lattice modules*, Fund. Math., **66** (1970), 341-371.

- [19] C. Jayaram and E. W. Johnson, *s-Prime elements in multiplicative lattices*, Period. Math. Hung., **31** (1995), 201-208.
- [20] F. Harary, *Graph Theory*, Narosa, New Delhi, 1988.
- [21] H. M. Nakkar, E. A. AL-Khouja and GH. Al-Mohammad, *A characterization of regular lattice modules by the radical elements*, 40th-science week at the University of Tachreen, **2** (2000), 63-75.
- [22] N. Phadatare, V. Kharat and S. Ballal, *Semi-complement graph of lattice modules*, Soft Computing, **23** (2019), 3973-3978.
- [23] S. Rajae, *Essential submodules relative to a submodule*, J. Algebra Relat. Topics, (2) **11** (2023), 59-71.
- [24] D. B. West, *Introduction to graph theory*, Second ed., Prentice-Hall of India, New Delhi, 2002.